



Available online at www.sciencedirect.com



JOURNAL OF Approximation Theory

Journal of Approximation Theory 253 (2020) 105378

www.elsevier.com/locate/jat

Full Length Article

# Asymptotics of the generalized Gegenbauer functions of fractional degree<sup>☆</sup>

Wenjie Liu<sup>a</sup>, Li-Lian Wang<sup>b,\*</sup>

<sup>a</sup> Department of Mathematics, Harbin Institute of Technology, Harbin, 150001, PR China <sup>b</sup> Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, 637371, Singapore

Received 5 April 2019; received in revised form 8 September 2019; accepted 30 January 2020 Available online 6 February 2020

Communicated by R. Wong

# Abstract

The generalized Gegenbauer functions of fractional degree (GGF-Fs), denoted by  ${}^{r}G_{\nu}^{(\lambda)}(x)$  (right GGF-Fs) and  ${}^{l}G_{\nu}^{(\lambda)}(x)$  (left GGF-Fs) with  $x \in (-1, 1)$ ,  $\lambda > -1/2$  and real  $\nu \ge 0$ , are special functions (usually non-polynomials), which are defined upon the hypergeometric representation of the classical Gegenbauer polynomial by allowing integer degree to be real fractional degree. Remarkably, the GGF-Fs become indispensable for optimal error estimates of polynomial approximation to singular functions, and have intimate relations with several families of nonstandard basis functions recently introduced for solving fractional differential equations. However, some properties of GGF-Fs, which are important pieces for the analysis and applications, are unknown or under explored. The purposes of this paper are twofold. The first is to show that for  $\lambda > -1/2$  and  $x = \cos \theta$  with  $\theta \in (0, \pi)$ ,

$$(\sin\theta)^{\lambda r} G_{\nu}^{(\lambda)}(\cos\theta) = \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi}(\nu + \lambda)^{\lambda}} \cos((\nu + \lambda)\theta - \lambda\pi/2) + \mathcal{R}_{\nu}^{(\lambda)}(\theta),$$

and derive the precise expression of the "residual" term  $\mathcal{R}_{\nu}^{(\lambda)}(\theta)$  for all real  $\nu \geq \nu_0$  (with some  $\nu_0 > 0$ ). With this at our disposal, we obtain the bounds of GGF-Fs uniform in  $\nu$ . Under an appropriate weight function, the bounds are uniform for  $\theta \in [0, \pi]$  as well. Moreover, we can study the asymptotics of

https://doi.org/10.1016/j.jat.2020.105378

0021-9045/© 2020 Elsevier Inc. All rights reserved.

 $<sup>\</sup>stackrel{\circ}{\sim}$  The research of the first author was supported by the China Postdoctoral Science Foundation Funded Project (No. 2017M620113), the National Natural Science Foundation of China (Nos. 11801120, 71773024 and 11271100), the Fundamental Research Funds for the Central Universities, China (Grant No. HIT.NSRIF.2019058 and No.HIT.NSRIF.2020081) and the Natural Science Foundation of Heilongjiang Province of China (Nos. A2016003 and G2018006). The research of the second author is partially supported by Singapore MOE AcRF Tier 2 Grants: MOE2018-T2-1-059 and MOE2017-T2-2-144.

<sup>\*</sup> Corresponding author.

E-mail addresses: liuwenjie@hit.edu.cn (W. Liu), lilian@ntu.edu.sg (L.-L. Wang).

GGF-Fs with large fractional degree  $\nu$ . The second is to present miscellaneous properties of GGF-Fs for better understanding of this family of useful special functions. (© 2020 Elsevier Inc. All rights reserved.

#### MSC: 30E15; 41A10; 41A25; 41A60; 65G50

*Keywords:* Generalized Gegenbauer functions of fractional degree; Asymptotic analysis; Riemann–Liouville fractional integrals and derivatives

#### 1. Introduction

Undoubtedly, polynomial approximation theory occupies a central place in algorithm development and numerical analysis of perhaps most of computational methods. Indeed, one finds numerous approximation results in various senses documented in a large volume of literature, which particularly include orthogonal polynomial approximation results related to spectral methods and hp-version finite element methods (see, e.g., [6,20,21,24] and the references therein). Typically, such results are established in Jacobi-weighted Sobolev spaces with integral-order regularity exponentials (see, e.g., [11,21]), or weighted Besov spaces with fractional regularity exponentials using the notion of space interpolation (see, e.g., [3-5]). In a very recent work [13], we introduced a new framework of fractional Sobolev-type spaces involving Riemann-Liouville (RL) fractional integrals and derivatives in the study of polynomial approximation to singular functions. Such spaces are naturally arisen from exact representations of orthogonal polynomial expansion coefficients, and could best characterize the fractional differentiability/regularity, leading to optimal error estimates. A very important piece of the puzzle therein is the so-called GGF-Fs that generalize the classical Gegenbauer polynomials of integer degree to functions of fractional degree. It is noteworthy that the GGF-Fs can be generalized by different means, e.g., the Rodrigues' formula and hypergeometric function representation. For instance, the right GGF-F:  ${}^{r}G_{u}^{(\lambda)}(x)$  can be viewed as special g-Jacobi functions (see Mirevski et al. [15]), defined by replacing the integer-order derivative in the Rodrigues' formula of the Jacobi polynomials by the RL fractional derivative. However, both the definition and derivation of some properties in [15] have flaws (see Remark 4.1). On the other hand, the Handbook [17, (15.9.15)] listed  ${}^{r}G_{\nu}^{(\lambda)}(x)$  but without presented any of their properties. Interestingly, as pointed out in [13], the GGF-Fs have a direct bearing on Jacobi polyfractonomials (cf. [28]) and generalized Jacobi functions (cf. [7,10]) recently introduced in developing efficient spectral methods for fractional differential equations. It is also noteworthy that the seminal work of Gui and Babuška [9] on hp-estimates of Legendre approximation of singular functions essentially relied on some non-classical Jacobi polynomials with the parameter  $\alpha$  or  $\beta < -1$ , which turned out closely related to GGF-Fs. In a nutshell, the GGF-Fs (and more generally the generalized Jacobi functions of fractional degree) can be of great value for numerical analysis and computational algorithms, but many of their properties are still under explored.

It is known that the study of asymptotics has been a longstanding subject of special functions and their far reaching applications (see, e.g., [16,17,23]). Most of the asymptotic results of classical orthogonal polynomials can be found in the books [17,22], and are reported in the review papers [14,26,27] in more general senses. We highlight that the asymptotic formulas of the hypergeometric function:  $_2F_1(a - \mu, b + \mu; c; (1 - z)/2)$  in terms of Bessel functions for large  $\mu$ , were derived in Jones [12] following the idea of Olver [16] using differential equations, where the representations with fewer restrictions on the parameters different from those in Watson [25] could be obtained. Farid Khwaja and Olde Daalhuis [8] discussed asymptotics of  ${}_{2}F_{1}(a - e_{1}\mu, b + e_{2}\mu; c + e_{3}\mu; (1 - z)/2)$  with  $e_{j} = 0, \pm 1, j = 1, 2, 3$  in terms of Bessel functions by using the contour integrals.

One of the main objectives of this paper is to derive the uniform bounds for the GGF-Fs, which are valid for real degree  $\nu > 0$  with fixed  $\lambda$ , and also for all  $\theta \in [0, \pi]$  but with a suitable weight function to absorb the singularities at the endpoints. As such, we can obtain the asymptotic formulas for large degree  $\nu$ , and some other useful estimates of the GGF-Fs. Our delicate analysis is based on an integral representation from a very useful fractional integral formula in [13] (see (2.7) and Lemma 2.1). In fact, the Watson's Lemma and asymptotic analysis for Legendre polynomials (cf. [16]) indeed cast light on our study. It is important to point out the GGF-Fs are defined as hypergeometric functions with special parameters (see Definition 2.1), so some asymptotic results follow from [8,12] for large parameters in terms of Bessel functions. However, we intend to derive the results uniform for the degree and the variable, and the estimates for large parameters are directly consequences. In other words, our study can lead to different and more explicitly informative estimates. As such, the results herein can offer useful tools for analysis of polynomial approximation and applications of this family of special functions. A second purpose of this paper is to present various properties of GGF-Fs. These particularly include singular behaviors of GGF-Fs in the vicinity of the endpoints, and useful fractional calculus formulas.

The paper is organized as follows. In Section 2, we first introduce the definition of GGF-Fs, and then present the main results. We then provide their proofs in Section 3. In Section 4, we present assorted properties of GGF-Fs for better understanding of this family of special functions.

#### 2. Main results

## 2.1. Generalized Gegenbauer functions of fractional degree

Different from Mirevski et al. [15], we follow [13] to define two types of GGF-Fs by the hypergeometric function.

**Definition 2.1.** For real  $\lambda > -1/2$ , we define the right GGF-F on (-1, 1) of real degree  $\nu \ge 0$  as

$${}^{r}G_{\nu}^{(\lambda)}(x) = {}_{2}F_{1}\left(-\nu,\nu+2\lambda;\lambda+\frac{1}{2};\frac{1-x}{2}\right) = 1 + \sum_{j=1}^{\infty} \frac{(-\nu)_{j}(\nu+2\lambda)_{j}}{j!(\lambda+1/2)_{j}} \left(\frac{1-x}{2}\right)^{j}, \quad (2.1)$$

and the left GGF-F of real degree  $\nu \ge 0$  as

$${}^{l}G_{\nu}^{(\lambda)}(x) = (-1)^{[\nu]} {}_{2}F_{1}\left(-\nu, \nu+2\lambda; \lambda+\frac{1}{2}; \frac{1+x}{2}\right) = (-1)^{[\nu]} {}^{r}G_{\nu}^{(\lambda)}(-x),$$
(2.2)

where  $[\nu]$  is the largest integer  $\leq \nu$ , and the Pochhammer symbol:  $(a)_j = a(a+1)\cdots(a+j-1)$ .

In the above, the hypergeometric function is a power series given by

$${}_{2}F_{1}(a,b;c;z) = 1 + \sum_{j=1}^{\infty} \frac{(a)_{j}(b)_{j}}{(c)_{j}} \frac{z^{j}}{j!},$$
(2.3)

where *a*, *b*, *c* are real, and  $-c \notin \mathbb{N} := \{1, 2, ...\}$  (see, e.g., [2]).

Note that if  $\nu = n \in \mathbb{N}_0 := \{0\} \cup \mathbb{N}$ , we have

$${}^{r}G_{n}^{(\lambda)}(x) = {}^{l}G_{n}^{(\lambda)}(x) = G_{n}^{(\lambda)}(x) = \frac{P_{n}^{(\lambda-1/2,\lambda-1/2)}(x)}{P_{n}^{(\lambda-1/2,\lambda-1/2)}(1)}, \quad \lambda > -\frac{1}{2},$$
(2.4)

where  $P_n^{(\alpha,\beta)}(x)$  is the classical Jacobi polynomial as defined in Szegö [22]. For  $\lambda = 1/2$ , the right GGF-F turns to be the Legendre function (cf. [23]):  ${}^rG_{\nu}^{(1/2)}(x) = P_{\nu}(x)$ . For  $\lambda = 0$ , we have

$${}^{r}G_{\nu}^{(0)}(x) = {}^{r}G_{\nu}^{(0)}(\cos\theta) = \cos(\nu\theta) = \cos(\nu \arccos x) := T_{\nu}(x), \tag{2.5}$$

thanks to the property (cf. [1, (15.1.17)]):

$$_{2}F_{1}(-a, a, 1/2; \sin^{2} t) = \cos(2at), \quad a, t \in \mathbb{R} := (-\infty, \infty).$$
 (2.6)

**Remark 2.1.** The GGF-Fs  ${}^{r}G_{n-\lambda+1/2}^{(\lambda)}(x)$  and  ${}^{l}G_{n-\lambda+1/2}^{(\lambda)}(x)$  with integer *n* up to some constant multiple, coincide with some nonstandard singular basis functions introduced in [7,28] for accurate solution of fractional differential equations.

Inherited from the Bateman's fractional integral formula for hypergeometric functions (cf. [2, P. 313]), we can derive the following very useful formula (cf. [13, Thm. 3.1]): for  $\lambda > -1/2$ , and real  $\nu \ge s \ge 0$ ,

$${}_{x}I_{1}^{s}\left\{(1-x^{2})^{\lambda-1/2} {}^{r}G_{\nu}^{(\lambda)}(x)\right\} = \frac{1}{\Gamma(s)} \int_{x}^{1} \frac{(1-y^{2})^{\lambda-1/2} {}^{r}G_{\nu}^{(\lambda)}(y)}{(y-x)^{1-s}} dy$$
  
=  $\frac{\Gamma(\lambda+1/2)}{2^{s}\Gamma(\lambda+s+1/2)} (1-x^{2})^{\lambda+s-1/2} {}^{r}G_{\nu-s}^{(\lambda+s)}(x),$  (2.7)

where  $_{x}I_{1}^{s}$  is the right-sided RL fractional integral operator defined by

$${}_{x}I_{1}^{s}u(x) = \frac{1}{\Gamma(s)} \int_{x}^{1} \frac{u(y)}{(y-x)^{1-s}} dy.$$
(2.8)

Note that a similar formula is available for the left GGF-F  ${}^{l}G_{\nu}^{(\lambda)}(x)$  but associated with the left-sided RL fractional integral.

Thanks to (2.7), we can derive the following formula crucial for the analysis.

**Lemma 2.1.** For real  $\nu, \lambda \ge 0$ , we have

$$(\sin\theta)^{2\lambda-1} {}^{r} G_{\nu}^{(\lambda)}(\cos\theta) = \frac{2^{\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi} \Gamma(\lambda)} \int_{0}^{\theta} \frac{\cos((\nu+\lambda)\phi)}{(\cos\phi-\cos\theta)^{1-\lambda}} d\phi,$$
(2.9)

for any  $\theta \in (0, \pi)$ .

**Proof.** From (2.5) and (2.7) with  $\lambda = 0$ , we obtain immediately that for  $\nu \ge s \ge 0$ ,

$$(1-x^2)^{s-1/2} {}^r G_{\nu-s}^{(s)}(x) = \frac{2^s \Gamma(s+1/2)}{\sqrt{\pi} \Gamma(s)} \int_x^1 \frac{1}{(y-x)^{1-s}} \frac{T_\nu(y)}{\sqrt{1-y^2}} \, dy.$$
(2.10)

Substituting s and v in the above identity by  $\lambda$  and  $\nu + \lambda$ , respectively, and using a change of variables:  $x = \cos \theta$  and  $y = \cos \phi$ , we derive (2.9) from (2.10) straightforwardly.

**Remark 2.2.** If  $\lambda = 1/2$  and  $\nu = n$ , the identity (2.9) leads to the first Dirichlet–Mehler formula for the Legendre polynomial (cf. [23, (6.51)]):

$$P_n(\cos\theta) = \frac{\sqrt{2}}{\pi} \int_0^\theta \frac{\cos(n+1/2)\phi}{\sqrt{\cos\phi - \cos\theta}} \, d\phi, \qquad \theta \in (0,\pi), \quad n \in \mathbb{N}_0.$$
(2.11)

One approach to obtain the asymptotic formula for Legendre polynomial with  $n \to \infty$  is based on this formula, and the Watson's lemma (cf. [16, P. 113]). This useful argument indeed sheds light on the study of GGF-Fs herein. However, we aim to study the behavior of GGF-Fs uniform for all  $\nu$ , so the route appears very different, delicate and more involved.

# 2.2. Main results

We first state the results, whose proofs are given in Section 3. Here, we only consider the right GGF-Fs, but in view of (2.2), similar results can be obtained for the left counterparts. Note from (2.5) that if  $\lambda = 0$ , then  $\mathcal{R}_{\nu}^{(0)}(\theta) \equiv 0$  in (2.12). Thus, in what follows, we exclude this trivial case and assume that  $\lambda \neq 0$ .

**Theorem 2.1.** If  $\lambda > -1/2$  and  $\lambda \neq 0$ , then for all  $\theta \in (0, \pi)$ , we have

$$(\sin\theta)^{\lambda r} G_{\nu}^{(\lambda)}(\cos\theta) = \frac{2^{\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda}} \cos((\nu+\lambda)\theta - \lambda\pi/2) + \mathcal{R}_{\nu}^{(\lambda)}(\theta), \qquad (2.12)$$

where the "residual" term  $\mathcal{R}_{\nu}^{(\lambda)}(\theta)$  with a representation is given by (3.33), and there holds

$$|\mathcal{R}_{\nu}^{(\lambda)}(\theta)| \le \mathcal{S}_{\nu}^{(\lambda)}(\theta), \quad \forall \theta \in (0,\pi).$$
(2.13)

Here, the bound  $S_{\nu}^{(\lambda)}(\theta)$  is given by

(i) for 
$$0 < \lambda \leq 2, \nu + \lambda > 1$$
 and  $\nu > 0$ ,  

$$\mathcal{S}_{\nu}^{(\lambda)}(\theta) = \frac{\lambda|\lambda - 1|2^{\lambda}\Gamma(\lambda + 1/2)}{\sqrt{\pi}(\nu + \lambda - 1)^{\lambda + 1}} \Big\{ |\cot \theta| + \frac{2}{3} \frac{\lambda + 1}{\nu + \lambda - 1} \Big\};$$
(2.14)

(ii) for  $\lambda > 2$ ,  $\nu > \lambda - 3$  and  $\nu > 0$ ,

$$S_{\nu}^{(\lambda)}(\theta) = \frac{\lambda(\lambda-1)2^{3\lambda/2}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+1)^{\lambda+1}} \left\{ |\cot\theta| + \frac{2}{3}\frac{\lambda+1}{\nu+1} + \frac{2^{2-\lambda}\Gamma(2\lambda-1)}{\Gamma(\lambda+1)}\frac{(\nu+1)^{\lambda+1}}{(\nu-\lambda+3)^{2\lambda-1}} |\cot\theta|^{\lambda-2} \left( |\cot\theta| + \frac{2}{3}\frac{2\lambda-1}{\nu-\lambda+3} \right) \right\};$$

$$(2.15)$$

(iii) for  $-1/2 < \lambda < 0$  and  $\nu \ge 2$ ,

$$S_{\nu}^{(\lambda)}(\theta) = \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi}(\nu + \lambda - 1)^{\lambda + 1}} \left\{ 2(\lambda + 1/2) |\cot \theta| + \frac{(\lambda + 1)^2}{\nu + \lambda} + (\lambda + 2)(\lambda + 1) \left( 1 + \frac{2}{\nu + \lambda - 1} - \frac{\lambda(\lambda + 2)}{(\nu + \lambda - 1)^2} \right) \left( |\cot \theta| + \frac{2}{3} \frac{\lambda + 3}{\nu + \lambda - 1} \right) \right\}.$$
(2.16)

With Theorem 2.1 at our disposal, we can obtain the bound of  $S_{\nu}^{(\lambda)}(\theta)$ , and characterize its explicit dependence of  $\theta$  and decay rate in  $\nu$ .

**Corollary 2.1.** If  $\lambda > -1/2$  and  $\lambda \neq 0$ , then we have

$$\left| (\sin \theta)^{\lambda r} G_{\nu}^{(\lambda)}(\cos \theta) - \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi} (\nu + \lambda)^{\lambda}} \cos((\nu + \lambda)\theta - \lambda\pi/2) \right| \le \frac{B_{\nu}^{(\lambda)}}{\nu^{\lambda + 1} \sin \theta},$$
(2.17)

where the constant  $B_{\nu}^{(\lambda)}$  is given by

(i) for 
$$0 < \lambda \le 2$$
 and  $\nu + \lambda > 1$ ,  

$$B_{\nu}^{(\lambda)} = \frac{\lambda |\lambda - 1| 2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi}} \frac{3\nu + 5\lambda - 1}{3(\nu + \lambda - 1)} \exp\left(\frac{1 - \lambda^2}{\nu + \lambda - 1}\right),$$
(2.18)

and the bound (2.17) holds for all  $\theta \in (0, \pi)$ ; (ii) for  $\lambda > 2$  and  $\nu > \lambda - 3$ , we have

$$B_{\nu}^{(\lambda)} = \frac{\lambda(\lambda-1)2^{3\lambda/2}\Gamma(\lambda+1/2)}{3\sqrt{\pi}} \left\{ \frac{3\nu+2\lambda+5}{\nu+1} + (c\pi)^{\lambda-2} \frac{\Gamma(2\lambda-1)}{\Gamma(\lambda+1)} \frac{3\nu+\lambda+7}{\nu-\lambda+3} \exp\left(\frac{(2\lambda-5)(\lambda+1)}{\nu-\lambda+3}\right) \right\},$$
(2.19)

and the bound (2.17) holds for all  $\theta \in [cv^{-1}, \pi - cv^{-1}]$  with c being a fixed positive constant;

(iii) for 
$$-1/2 < \lambda < 0$$
 and  $\nu \ge 2$ ,

$$B_{\nu}^{(\lambda)} = \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi}} \frac{\nu^{\lambda+1}}{(\nu + \lambda - 1)^{\lambda+1}} \left\{ 2(\lambda + 1/2) + \frac{(\lambda + 1)^2}{\nu + \lambda} + (\lambda + 2)(\lambda + 1) \left( 1 + \frac{2}{\nu + \lambda - 1} - \frac{\lambda(\lambda + 2)}{(\nu + \lambda - 1)^2} \right) \left( 1 + \frac{2}{3} \frac{\lambda + 3}{\nu + \lambda - 1} \right) \right\},$$
(2.20)

and the bound (2.17) holds for all  $\theta \in (0, \pi)$ .

We provide the derivation of the above bounds right after the proof of Theorem 2.1. Note that in the second case:  $\lambda > 2$ , the bound is only available for  $\theta \in [cv^{-1}, \pi - cv^{-1}]$  with some fixed constant c > 0. Indeed, the situation is reminiscent to the classical Gegenbauer polynomial with asymptotics only valid for  $\theta \in [cn^{-1}, \pi - cn^{-1}]$  with large *n*, as we remark below.

**Remark 2.3.** From (2.4) and Theorem 2.1, we obtain that for  $\nu = n \in \mathbb{N}$ ,

$$(\sin\theta)^{\lambda} P_{n}^{(\lambda-1/2,\lambda-1/2)}(\cos\theta) = (\sin\theta)^{\lambda} P_{n}^{(\lambda-1/2,\lambda-1/2)}(1) G_{n}^{(\lambda)}(\cos\theta)$$
$$= \frac{2^{\lambda} \Gamma(n+\lambda+1/2)}{\sqrt{\pi} n! (n+\lambda)^{\lambda}} \cos((n+\lambda)\theta - \lambda\pi/2) + \frac{\Gamma(n+\lambda+1/2)}{\Gamma(\lambda+1/2)n!} \mathcal{R}_{n}^{(\lambda)}(\theta).$$
(2.21)

Then from Corollary 2.1, we can derive the bounds uniform for n. In fact, we can recover the asymptotic formula for the classical Gegenbauer polynomial with large n (cf. [22, Thm 8.21.13]):

$$(\sin\theta)^{\lambda} P_n^{(\lambda-1/2,\lambda-1/2)}(\cos\theta) = \frac{2^{\lambda}}{\sqrt{\pi n}} \{ \cos((n+\lambda)\theta - \lambda\pi/2) + (n\sin\theta)^{-1} O(1) \}, \quad (2.22)$$

for all  $\lambda > -1/2$ ,  $\lambda \neq 0$  and  $\theta \in [cn^{-1}, \pi - cn^{-1}]$  with  $n \gg 1$  and *c* being a fixed positive constant. Indeed, using the property of the Gamma function (cf. [1, (6.1.38)]):

$$\Gamma(x+1) = \sqrt{2\pi} x^{x+1/2} \exp\left(-x + \frac{\eta}{12x}\right), \quad x > 0, \quad 0 < \eta < 1,$$
(2.23)

and the bounds of  $\mathcal{R}_n^{(\lambda)}(\theta)$  in Corollary 2.1, we can deduce (2.22) straightforwardly.

Thanks to Theorem 2.1, we can derive the following uniform bounds for  $\theta \in [0, \pi]$ , and nearly all fractional degree  $\nu > 0$ . We refer to Section 3.4 for its proof.

**Theorem 2.2.** (i) If  $-1/2 < \lambda < 0$  and  $\nu \ge 2$ ; or  $0 < \lambda \le 2$ ,  $\nu + \lambda > 1$  and  $\nu > 0$ , then we have

$$\begin{aligned} |\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)| &= \left| (\sin \theta)^{\lambda+1} \, {}^{r} G_{\nu}^{(\lambda)}(\cos \theta) - \frac{2^{\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi} (\nu+\lambda)^{\lambda}} \, (\sin \theta) \cos((\nu+\lambda)\theta - \lambda\pi/2) \right| \\ &\leq \widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta), \end{aligned} \tag{2.24}$$

for all  $\theta \in [0, \pi]$ , where  $\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = (\sin \theta) \mathcal{R}_{\nu}^{(\lambda)}(\theta)$  and  $\widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta) = (\sin \theta) \mathcal{S}_{\nu}^{(\lambda)}(\theta)$ . (ii) If  $\lambda > 2$ ,  $\nu > \lambda - 3$  and  $\nu > 0$ , then we have

$$\begin{aligned} |\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)| &= \left| (\sin \theta)^{2\lambda - 1} \, {}^{r} G_{\nu}^{(\lambda)}(\cos \theta) - \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi} (\nu + \lambda)^{\lambda}} \, (\sin \theta)^{\lambda - 1} \cos((\nu + \lambda)\theta - \lambda \pi/2) \right| \\ &\leq \widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta), \end{aligned}$$

$$(2.25)$$

for all  $\theta \in [0, \pi]$ , where

$$\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = (\sin \theta)^{\lambda - 1} \, \mathcal{R}_{\nu}^{(\lambda)}(\theta), \quad \widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta) = (\sin \theta)^{\lambda - 1} \, \mathcal{S}_{\nu}^{(\lambda)}(\theta).$$

In the end of this section, we provide some numerical illustrations of the uniform bounds in Theorem 2.2. In Fig. 2.1, we plot the graphs of  $|\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)|$  and  $\widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta)$  for  $\theta \in [0, \pi]$  and with  $\lambda = -0.4, -0.2, 0.7, 1.6, 2.3, 3.1, \nu = 20.3$ . Indeed, we observe that in all cases, the curves of the upper bounds are on the top of  $|\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)|$ , and the "sharp" corner of  $\widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta)$  at  $\theta = \pi/2$  is largely due to the involved  $|\cos \theta|$ .

### 3. Proofs of the main results

#### 3.1. Some lemmas

As the proofs of the main results are quite involved, we take several steps and summarize the intermediate results in three lemmas.

**Lemma 3.1.** For real  $\lambda > 0$ ,  $\theta \in (0, \pi)$  and t > 0, define

$$g(\theta, t) \coloneqq \frac{\cos(\theta - it) - \cos\theta}{t} = \frac{\cos\theta \left(\cosh t - 1\right) + i\sin\theta \sinh t}{t},$$
  

$$f^{(\lambda)}(\theta, t) \coloneqq \frac{g^{\lambda - 1}(\theta, t) - g^{\lambda - 1}(\theta, 0)}{t}, \quad g(\theta, 0) \coloneqq \lim_{t \to 0^+} g(\theta, t) = i\sin\theta.$$
(3.1)

Then we have for  $\theta \in (0, \pi)$  and t > 0,

(i) for  $0 < \lambda \leq 2$ ,

$$|f^{(\lambda)}(\theta,t)| \le |\lambda - 1| (\sin \theta)^{\lambda - 1} \left( |\cot \theta| + \frac{2t}{3} \right) e^t;$$
(3.2)

(ii) for  $\lambda > 2$ ,

$$|f^{(\lambda)}(\theta,t)| \le 2^{\lambda/2} (\lambda-1) (\sin\theta)^{\lambda-1} \left( |\cot\theta| + \frac{2t}{3} \right) \left( 1 + \frac{|\cot\theta|^{\lambda-2}}{2^{\lambda-2}} t^{\lambda-2} e^{(\lambda-2)t} \right) e^{(\lambda-1)t}.$$
(3.3)



Fig. 2.1. Plots of  $|\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)|$  and  $\widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta)$  in Theorem 2.2, where  $\theta \in [0, \pi], \lambda = -0.4, -0.2, 0.7, 1.6, 2.3, 3.1$  and  $\nu = 20.3$ .

To avoid distracting from proving the main result, we put this a bit lengthy proof but only involving fundamental calculus in Appendix A.

A critical step is to show that the integral in (2.9) satisfies the following identity.

**Lemma 3.2.** For real  $\nu, \lambda \ge 0$ , and  $\theta \in (0, \pi)$ , we have

$$\int_{0}^{\theta} \frac{\cos((\nu+\lambda)\phi)}{(\cos\phi-\cos\theta)^{1-\lambda}} d\phi = \frac{\Gamma(\lambda)}{(\nu+\lambda)^{\lambda}} \frac{\cos((\nu+\lambda)\theta-\lambda\pi/2)}{(\sin\theta)^{1-\lambda}} + \check{\mathcal{R}}_{\nu}^{(\lambda)}(\theta), \tag{3.4}$$

where

$$\check{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) \coloneqq \int_{0}^{\infty} \Re \left\{ i \, e^{-i(\nu+\lambda)\theta} \, f^{(\lambda)}(\theta, t) \right\} t^{\lambda} e^{-(\nu+\lambda)t} dt, \tag{3.5}$$

and  $f^{(\lambda)}(\theta, t)$  is defined in (3.1).



**Fig. 3.2.** Contour integral for (3.8). Left: for  $\lambda \ge 1$ ; Right: for  $0 < \lambda < 1$ .

**Proof.** It is evident that by the parity, we have

$$\int_0^\theta \frac{\cos((\nu+\lambda)\phi)}{(\cos\phi - \cos\theta)^{1-\lambda}} \, d\phi = \frac{1}{2} \int_{-\theta}^\theta F_\nu^{(\lambda)}(\theta,\phi) \, d\phi \,, \tag{3.6}$$

where we denote

$$F_{\nu}^{(\lambda)}(\theta,\phi) \coloneqq \frac{e^{\mathrm{i}(\nu+\lambda)\phi}}{(\cos\phi - \cos\theta)^{1-\lambda}} \,. \tag{3.7}$$

We consider the cases with  $\lambda \ge 1$  and  $0 < \lambda < 1$ , separately.

(i) *Proof of* (3.4) *with*  $\lambda \ge 1$ . From the Cauchy–Goursat theorem, we infer that for any fixed  $\theta \in (0, \pi)$  and real  $\nu > 0$ , the contour integration of  $F_{\nu}^{(\lambda)}(\theta, \cdot)$  (with an extension to the complex plane) along the rectangular contour in Fig. 3.2 (left), is zero. Thus, we have

$$\int_{-\theta}^{\theta} F_{\nu}^{(\lambda)}(\theta,\phi) d\phi = \int_{-\theta}^{-\theta+iR} F_{\nu}^{(\lambda)}(\theta,\phi) d\phi - \int_{\theta}^{\theta+iR} F_{\nu}^{(\lambda)}(\theta,\phi) d\phi + \int_{-\theta+iR}^{\theta+iR} F_{\nu}^{(\lambda)}(\theta,\phi) d\phi = i \int_{0}^{R} \left\{ F_{\nu}^{(\lambda)}(\theta,-\theta+it) - F_{\nu}^{(\lambda)}(\theta,\theta+it) \right\} dt + \int_{-\theta}^{\theta} F_{\nu}^{(\lambda)}(\theta,t+iR) dt,$$
(3.8)

where we made the change of variables for three integrals:  $\phi = -\theta + it, \theta + it, t + iR$ , respectively.

For  $\lambda \ge 1$  and R > 0, we have

$$|F_{\nu}^{(\lambda)}(\theta, t + iR)| = \frac{|e^{i(\nu+\lambda)t - (\nu+\lambda)R}|}{|\cos(t + iR) - \cos\theta|^{1-\lambda}}$$
  
=  $e^{-(\nu+\lambda)R} ((\cos t \cosh R - \cos\theta)^2 + \sin^2 t \sinh^2 R)^{(\lambda-1)/2}$   
 $\leq e^{-(\nu+\lambda)R} ((\cosh R + 1)^2 + \sinh^2 R)^{(\lambda-1)/2}$   
=  $2^{(1-\lambda)/2} e^{-(\nu+1)R} (1 + 2e^{-R} + 2e^{-2R} + 2e^{-3R} + e^{-4R})^{(\lambda-1)/2}$   
 $< 2^{\lambda-1} e^{-(\nu+1)R}.$  (3.9)

Thus, we have

$$\lim_{R \to \infty} \int_{-\theta + iR}^{\theta + iR} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi = \lim_{R \to \infty} \int_{-\theta}^{\theta} F_{\nu}^{(\lambda)}(\theta, t + iR) dt = 0.$$
(3.10)

Recall the notation in (3.1):  $tg(\theta, t) = \cos(\theta - it) - \cos\theta$ . In view of (3.1), we can write  $g^{\lambda-1}(\theta, t) = g^{\lambda-1}(\theta, 0) + tf^{(\lambda)}(\theta, t)$ . Thus, by a direct calculation, we obtain

$$iF_{\nu}^{(\lambda)}(\theta, -\theta + it) - iF_{\nu}^{(\lambda)}(\theta, \theta + it) = \{ig^{\lambda-1}(\theta, t)e^{-i(\nu+\lambda)\theta} + (ig^{\lambda-1}(\theta, t)e^{-i(\nu+\lambda)\theta})^*\}t^{\lambda-1}e^{-(\nu+\lambda)t}$$

$$= 2\Re\{ig^{\lambda-1}(\theta, t)e^{-i(\nu+\lambda)\theta}\}t^{\lambda-1}e^{-(\nu+\lambda)t}$$

$$= 2\Re\{ig^{\lambda-1}(\theta, 0)e^{-i(\nu+\lambda)\theta}\}t^{\lambda-1}e^{-(\nu+\lambda)t}$$

$$+ 2\Re\{if^{(\lambda)}(\theta, t)e^{-i(\nu+\lambda)\theta}\}t^{\lambda}e^{-(\nu+\lambda)t}.$$
(3.11)

Since  $i = e^{i\pi/2}$  and  $g(\theta, 0) = (i \sin \theta)^{\lambda-1}$ , we have

$$\begin{aligned} \Re \left\{ i g^{\lambda - 1}(\theta, 0) e^{-i(\nu + \lambda)\theta} \right\} &= \Re \left\{ i (i \sin \theta)^{\lambda - 1} e^{-i(\nu + \lambda)\theta} \right\} \\ &= (\sin \theta)^{\lambda - 1} \Re \left\{ e^{-i((\nu + \lambda)\theta - \lambda\pi/2)} \right\} \\ &= (\sin \theta)^{\lambda - 1} \cos((\nu + \lambda)\theta - \lambda\pi/2). \end{aligned}$$
(3.12)

Using the definition of the Gamma function, we find that for any a > 0, z > -1,

$$\int_0^\infty t^z e^{-at} dt = \frac{\Gamma(z+1)}{a^{z+1}}, \quad \text{as} \quad \Gamma(z+1) = \int_0^\infty e^{-t} t^z dt.$$
(3.13)

As a direct consequence of (3.12)–(3.13), we have

$$2\int_0^\infty \Re\left\{i\,g^{\lambda-1}(\theta,0)e^{-i(\nu+\lambda)\theta}\right\}t^{\lambda-1}e^{-(\nu+\lambda)t}dt = \frac{2\,\Gamma(\lambda)}{(\nu+\lambda)^\lambda}\,\frac{\cos((\nu+\lambda)\theta-\lambda\pi/2)}{(\sin\theta)^{1-\lambda}}.$$
(3.14)

Letting  $R \to \infty$  in (3.8), we obtain (3.4)–(3.5) from (3.6), (3.10)–(3.11) and (3.14) directly.

(ii) Proof of (3.4) with  $0 < \lambda < 1$ . In this case, we integrate along a similar contour but exclude singular points  $\phi = \pm \theta$ , as depicted in Fig. 3.2 (right), where  $0 < \epsilon < \theta$ . Like (3.8), we have

$$\int_{-\theta+\epsilon}^{\theta-\epsilon} F_{\nu}^{(\lambda)}(\theta,\phi) d\phi = I_1(\epsilon,R) + I_2(R) + I_3(\epsilon) + I_4(\epsilon), \qquad (3.15)$$

where

$$I_{1}(\epsilon, R) := \int_{-\theta+i\epsilon}^{-\theta+iR} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi - \int_{\theta+i\epsilon}^{\theta+iR} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi,$$

$$I_{2}(R) := \int_{-\theta+iR}^{\theta+iR} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi,$$

$$I_{3}(\epsilon) := \int_{-\theta+\epsilon}^{-\theta+\epsilon+i\epsilon} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi - \int_{\theta-\epsilon}^{\theta-\epsilon+i\epsilon} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi$$

$$I_{4}(\epsilon) := -\int_{-\theta+i\epsilon}^{-\theta+\epsilon+i\epsilon} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi - \int_{\theta-\epsilon+i\epsilon}^{\theta+i\epsilon} F_{\nu}^{(\lambda)}(\theta, \phi) d\phi.$$
(3.16)

Using a change of variable:  $\phi = \pm \theta + it$ , and noting that the derivation in (3.11)–(3.12) is valid for  $0 < \lambda < 1$ , we have

$$I_{1}(\epsilon, R) = i \int_{\epsilon}^{R} \left\{ F_{\nu}^{(\lambda)}(\theta, -\theta + it) - F_{\nu}^{(\lambda)}(\theta, \theta + it) \right\} dt$$
  
$$= 2 \frac{\cos((\nu + \lambda)\theta - \lambda\pi/2)}{(\sin\theta)^{1-\lambda}} \int_{\epsilon}^{R} t^{\lambda-1} e^{-(\nu+\lambda)t} dt$$
  
$$+ 2 \int_{\epsilon}^{R} \Re \left\{ i f^{(\lambda)}(\theta, t) e^{-i(\nu+\lambda)\theta} \right\} t^{\lambda} e^{-(\nu+\lambda)t} dt.$$
  
(3.17)

From (3.2) and (3.13)-(3.14), we infer that

$$\lim_{\epsilon \to 0; R \to \infty} I_1(\epsilon, R) = \frac{2 \Gamma(\lambda)}{(\nu + \lambda)^{\lambda}} \frac{\cos((\nu + \lambda)\theta - \lambda\pi/2)}{(\sin \theta)^{1 - \lambda}} + 2 \int_0^\infty \Re\{i f^{(\lambda)}(\theta, t)e^{-i(\nu + \lambda)\theta}\}t^{\lambda}e^{-(\nu + \lambda)t}dt.$$
(3.18)

Therefore, it suffices to show

$$\lim_{R \to \infty} I_2(R) = 0, \quad \lim_{\epsilon \to 0} I_3(\epsilon) = \lim_{\epsilon \to 0} I_4(\epsilon) = 0.$$
(3.19)

By (3.8), we have

$$I_2(R) = \int_{-\theta}^{\theta} F_{\nu}^{(\lambda)}(\theta, t + iR) dt, \qquad (3.20)$$

and

$$|F_{\nu}^{(\lambda)}(\theta, t + iR)| = \frac{|e^{i(\nu+\lambda)t - (\nu+\lambda)R}|}{|\cos(t + iR) - \cos\theta|^{1-\lambda}}$$
  
=  $e^{-(\nu+\lambda)R} ((\cos t \cosh R - \cos\theta)^2 + \sin^2 t \sinh^2 R)^{(\lambda-1)/2}$   
 $\leq e^{-(\nu+\lambda)R} (\sinh R)^{\lambda-1} |\sin t|^{\lambda-1}.$  (3.21)

Thus, for  $0 < \lambda < 1$  and  $\theta \in (0, \pi)$ ,

$$|I_2(R)| \le \frac{2e^{-(\nu+\lambda)R}}{(\sinh R)^{1-\lambda}} \int_0^\theta \frac{1}{(\sin t)^{1-\lambda}} dt \to 0, \quad \text{as } R \to \infty.$$
(3.22)

Next, using a change of variable:  $\theta = -\theta + \epsilon + it$ ,  $\theta - \epsilon + it$ , respectively, for two integrals, we obtain from a direct calculation that

$$I_{3}(\epsilon) = i \int_{0}^{\epsilon} \left\{ F_{\nu}^{(\lambda)}(\theta, -\theta + \epsilon + it) - F_{\nu}^{(\lambda)}(\theta, \theta - \epsilon + it) \right\} dt$$
  
$$= 2 \int_{0}^{\epsilon} \Re \left\{ \frac{i e^{-i(\nu + \lambda)(\theta - \epsilon)}}{(\cos(\theta - \epsilon - it) - \cos\theta)^{1 - \lambda}} \right\} e^{-(\nu + \lambda)t} dt.$$
(3.23)

Note that we have

$$|\cos(\theta - \epsilon - it) - \cos\theta| = ((\cos(\theta - \epsilon)\cosh t - \cos\theta)^2 + \sin^2(\theta - \epsilon)\sinh^2 t)^{1/2}$$
  

$$\geq |\sin(\theta - \epsilon)||\sinh t| \geq |\sin(\theta - \epsilon)||\sin t|,$$
(3.24)

where we used the inequality:  $|\sin t| \le \sinh t$  for t > 0 (cf. [17, (4.18.9)]). Therefore, for  $0 < \lambda < 1$ , we have

$$|I_3(\epsilon)| \le \frac{2}{(\sin(\theta - \epsilon))^{1-\lambda}} \int_0^{\epsilon} (\sin t)^{\lambda - 1} dt \to 0, \quad \text{as } \epsilon \to 0.$$
(3.25)

Similarly, with a change of variable:  $\theta = -\theta + i\epsilon + t$ ,  $\theta + i\epsilon - t$ , respectively, for two integrals, we derive

$$I_{4}(\epsilon) = -\int_{0}^{\epsilon} \left\{ F_{\nu}^{(\lambda)}(\theta, -\theta + i\epsilon + t) + F_{\nu}^{(\lambda)}(\theta, \theta + i\epsilon - t) \right\} dt$$
  
$$= -2e^{-\epsilon(\nu+\lambda)} \int_{0}^{\epsilon} \Re \left\{ \frac{e^{i(\nu+\lambda)(t-\theta)}}{(\cos(t-\theta + i\epsilon) - \cos\theta)^{1-\lambda}} \right\} dt.$$
(3.26)

It is evident that

$$|\cos(t - \theta + i\epsilon) - \cos\theta| = ((\cos(t - \theta)\cosh\epsilon - \cos\theta)^{2} + \sin^{2}(t - \theta)\sinh^{2}\epsilon)^{1/2} \ge \sin(\theta - t)\sinh\epsilon,$$
(3.27)

where as  $0 < \theta < \pi$  and  $0 < t < \epsilon < \theta$ , we have

 $0 < \theta - \epsilon < \theta - t < \theta < \pi.$ 

Using the fundamental inequalities:

$$\frac{2}{\pi}z \le \sin z \le z, \quad z \in (0, \pi/2), \tag{3.28}$$

we find

$$\frac{1}{\sin z} = \frac{1}{\sin(\pi - z)} \le \frac{\pi}{2} \max\left\{\frac{1}{z}, \frac{1}{\pi - z}\right\}, \quad z \in (0, \pi).$$
(3.29)

This implies

$$\sin(\theta - t) \ge \frac{2}{\pi} \min\{\theta - t, \pi - \theta + t\} > \frac{2}{\pi} \min\{\theta - \epsilon, \pi - \theta\}.$$
(3.30)

From (3.26) and (3.30), we obtain

$$|I_4(\epsilon)| \le 2e^{-\epsilon(\nu+\lambda)} \int_0^{\epsilon} |\cos(t-\theta+i\epsilon)-\cos\theta|^{\lambda-1} dt$$
  
$$\le \frac{2^{\lambda}}{\pi^{\lambda-1}} \frac{\epsilon}{(\sinh\epsilon)^{1-\lambda}} \max\{(\theta-\epsilon)^{\lambda-1}, (\pi-\theta)^{\lambda-1}\} \to 0, \text{ as } \epsilon \to 0.$$
(3.31)

Thus, letting  $\epsilon \to 0$  and  $R \to \infty$  in (3.15), we obtain (3.4)–(3.5) with  $0 < \lambda < 1$  from (3.6), (3.18), (3.22), (3.25) and (3.31).

In order to get a uniform coefficient in front of the cosine in (2.12) of Theorem 2.1 for the case  $-\frac{1}{2} < \lambda < 0$ , we need to use the following bound, which can be derived from [13, Thms 2.1-2.2] (see Proposition 4.7), and whose derivation is sketched in Appendix B.

**Lemma 3.3.** For  $0 < \lambda < 1$  and real  $v \ge 0$ , we have

$$\max_{|x| \le 1} \left\{ (1 - x^2)^{\lambda/2} \left|^r G_{\nu}^{(\lambda)}(x) \right| \right\} \le \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi} (\nu + \lambda)^{\lambda}}.$$
(3.32)

# 3.2. Proof of Theorem 2.1

Equipped with Lemmas 3.1–3.3, we are ready to prove the main results stated in Theorem 2.1. We first show Case-(i) and Case-(ii), as the proof of Case-(iii) needs to use the results of these two cases.

From (2.9) and Lemma 3.2, we derive

$$\mathcal{R}_{\nu}^{(\lambda)}(\theta) = \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi} \Gamma(\lambda)} (\sin \theta)^{1-\lambda} \check{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = \frac{2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi} \Gamma(\lambda)} (\sin \theta)^{1-\lambda} \int_{0}^{\infty} \Re \{ i e^{-i(\nu+\lambda)\theta} f^{(\lambda)}(\theta, t) \} t^{\lambda} e^{-(\nu+\lambda)t} dt.$$
(3.33)

We now estimate  $\check{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)$  in (3.4)–(3.5) by using Lemma 3.1. (i) For  $0 < \lambda \leq 2$  and  $\nu + \lambda > 1$ , we obtain from (3.2) and (3.13) that

$$\begin{aligned} |\breve{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)| &\leq \int_{0}^{\infty} |f^{(\lambda)}(\theta, t)| t^{\lambda} e^{-(\nu+\lambda)t} dt \\ &\leq |\lambda - 1| (\sin \theta)^{\lambda - 1} \int_{0}^{\infty} \left( |\cot \theta| + \frac{2t}{3} \right) t^{\lambda} e^{-(\nu+\lambda - 1)t} dt \\ &= \frac{|\lambda - 1| \Gamma(\lambda + 1)}{(\nu+\lambda - 1)^{\lambda+1}} (\sin \theta)^{\lambda - 1} \left( |\cot \theta| + \frac{2}{3} \frac{\lambda + 1}{\nu+\lambda - 1} \right). \end{aligned}$$
(3.34)

(ii) For  $\lambda > 2$  and  $\nu > \lambda - 3$ , we derive from (3.3) and (3.13) that

$$\begin{split} |\breve{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)| &\leq \int_{0}^{\infty} |f^{(\lambda)}(\theta,t)| t^{\lambda} e^{-(\nu+\lambda)t} dt &\leq 2^{\lambda/2} (\lambda-1) (\sin\theta)^{\lambda-1} \\ &\times \int_{0}^{\infty} \left( |\cot\theta| + \frac{2t}{3} \right) \left( 1 + \frac{|\cot\theta|^{\lambda-2}}{2^{\lambda-2}} t^{\lambda-2} e^{(\lambda-2)t} \right) t^{\lambda} e^{-(\nu+1)t} dt \\ &= \frac{2^{\lambda/2} (\lambda-1) \Gamma(\lambda+1)}{(\nu+1)^{\lambda+1}} (\sin\theta)^{\lambda-1} \left\{ |\cot\theta| + \frac{2}{3} \frac{\lambda+1}{\nu+1} \right. \\ &+ \frac{2^{2-\lambda} \Gamma(2\lambda-1)}{\Gamma(\lambda+1)} \frac{(\nu+1)^{\lambda+1}}{(\nu-\lambda+3)^{2\lambda-1}} |\cot\theta|^{\lambda-2} \left( |\cot\theta| + \frac{2}{3} \frac{2\lambda-1}{\nu-\lambda+3} \right) \right\}. \end{split}$$
(3.35)

Thus, we can derive the bounds in (2.14)–(2.15) from (3.33) and (3.34)–(3.35), respectively. (iii) We now turn to the case with  $-\frac{1}{2} < \lambda < 0$ , and start with the identity

$${}^{r}G_{\nu}^{(\lambda)}(x) = x \, {}^{r}G_{\nu-1}^{(\lambda+1)}(x) - \frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)}(1-x^{2}) \, {}^{r}G_{\nu-2}^{(\lambda+2)}(x), \quad \nu \ge 2, \tag{3.36}$$

which can be derived from (2.1) and the property of hypergeometric functions (see Proposition 4.5). Taking  $x = \cos\theta$  in (3.36) and multiplying the resulted identity by  $(\sin\theta)^{\lambda}$ , yields

$$(\sin\theta)^{\lambda r} G_{\nu}^{(\lambda)}(\cos\theta) = (\sin\theta)^{\lambda} \cos\theta^{r} G_{\nu-1}^{(\lambda+1)}(\cos\theta) - \frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)} (\sin\theta)^{\lambda+2r} G_{\nu-2}^{(\lambda+2)}(\cos\theta).$$
(3.37)

Thus, we have

$$\mathcal{R}_{\nu}^{(\lambda)}(\theta) = (\sin\theta)^{\lambda} \cos\theta^{r} G_{\nu-1}^{(\lambda+1)}(\cos\theta) + \widehat{\mathcal{R}}_{\nu}^{(\lambda)}(\theta), \qquad (3.38)$$

where

$$\widehat{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) \coloneqq -\frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)}(\sin\theta)^{\lambda+2} {}^{r}G_{\nu-2}^{(\lambda+2)}(\cos\theta) -\frac{2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda}}\cos((\nu+\lambda)\theta-\lambda\pi/2).$$

Using Lemma 3.3 with  $\lambda + 1 \in (1/2, 1)$ , we can bound the first term in (3.38) by

$$\begin{aligned} \left| (\sin \theta)^{\lambda} \cos \theta^{r} G_{\nu-1}^{(\lambda+1)} (\cos \theta) \right| &= \left| \cot \theta \left( \sin \theta \right)^{\lambda+1} {}^{r} G_{\nu-1}^{(\lambda+1)} (\cos \theta) \right| \\ &\leq \frac{2^{\lambda+1} \Gamma(\lambda+3/2)}{\sqrt{\pi} (\nu+\lambda)^{\lambda+1}} \left| \cot \theta \right|. \end{aligned}$$
(3.39)

Now, we deal with  $\widehat{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)$ . Using the proved result (2.12) with  $\lambda + 2 \in (3/2, 2)$ , we obtain that for  $\nu \geq 2$ ,

$$(\sin\theta)^{\lambda+2} {}^{r} G_{\nu-2}^{(\lambda+2)}(\cos\theta) = -\frac{2^{\lambda+2} \Gamma(\lambda+5/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda+2}} \cos((\nu+\lambda)\theta - \lambda\pi/2) + \mathcal{R}_{\nu-2}^{(\lambda+2)}(\theta).$$

Inserting it into  $\widehat{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)$  leads to

$$\widehat{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = \left\{ \frac{2^{\lambda} \Gamma(\lambda+1/2)(\nu-1)(\nu+2\lambda+1)}{\sqrt{\pi}(\nu+\lambda)^{\lambda+2}} - \frac{2^{\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda}} \right\} \cos((\nu+\lambda)\theta - \lambda\pi/2) \quad (3.40) \\
- \frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)} \mathcal{R}_{\nu-2}^{(\lambda+2)}(\theta).$$

Direct calculation yields

$$\left|\frac{2^{\lambda}\Gamma(\lambda+1/2)(\nu-1)(\nu+2\lambda+1)}{\sqrt{\pi}(\nu+\lambda)^{\lambda+2}} - \frac{2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda}}\right| \left|\cos((\nu+\lambda)\theta - \lambda\pi/2)\right| \\
\leq \frac{2^{\lambda}(\lambda+1)^{2}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda+2}}.$$
(3.41)

Using the proved bound in (2.14), we find

$$\frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)} \left| \mathcal{R}_{\nu-2}^{(\lambda+2)}(\theta) \right| \leq \frac{(\lambda+2)(\lambda+1)2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda-1)^{\lambda+1}} \\ \times \left\{ 1 + \frac{2}{\nu+\lambda-1} - \frac{\lambda(\lambda+2)}{(\nu+\lambda-1)^2} \right\} \left\{ |\cot\theta| + \frac{2}{3}\frac{\lambda+3}{\nu+\lambda-1} \right\}.$$
(3.42)

In view of (3.41)–(3.42), we obtain from (3.40) that

$$\begin{aligned} \left|\widehat{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)\right| &\leq \frac{2^{\lambda}(\lambda+1)^{2}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda+2}} + \frac{(\lambda+2)(\lambda+1)2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda-1)^{\lambda+1}} \\ &\times \left\{1 + \frac{2}{\nu+\lambda-1} - \frac{\lambda(\lambda+2)}{(\nu+\lambda-1)^{2}}\right\} \left\{|\cot\theta| + \frac{2}{3}\frac{\lambda+3}{\nu+\lambda-1}\right\}. \end{aligned}$$
(3.43)

Thanks to (3.39) and (3.43), we obtain from (3.38) the bound (2.16). This completes the proof of Theorem 2.1.

# 3.3. Proof of Corollary 2.1

We prove three cases separately.

(i) We obtain from (2.14) that

$$\begin{split} \nu^{\lambda+1} \sin \theta \, |\mathcal{R}_{\nu}^{(\lambda)}(\theta)| \\ &\leq \frac{\lambda|\lambda-1|2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}} \Big(|\cos \theta| + \frac{2}{3} \frac{(\lambda+1)\sin \theta}{\nu+\lambda-1} \Big) \Big(1 + \frac{1-\lambda}{\nu+\lambda-1} \Big)^{\lambda+1} \\ &\leq \frac{\lambda|\lambda-1|2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}} \Big(1 + \frac{2}{3} \frac{\lambda+1}{\nu+\lambda-1} \Big) \Big(1 + \frac{1-\lambda}{\nu+\lambda-1} \Big)^{\lambda+1}. \end{split}$$

Using the basic inequality:  $\ln(1 + z) \le z$  for z > -1, we find

$$\left(1+\frac{1-\lambda}{\nu+\lambda-1}\right)^{\lambda+1} = \exp\left((\lambda+1)\ln\left(1+\frac{1-\lambda}{\nu+\lambda-1}\right)\right) \le \exp\left(\frac{1-\lambda^2}{\nu+\lambda-1}\right).$$

Thus, we obtain  $B_{\nu}^{(\lambda)}$  immediately from the above for this case.

(ii) For  $\lambda > 2$ ,  $\nu - \lambda + 3 \ge 0$  and  $\theta \in [c\nu^{-1}, \pi - c\nu^{-1}]$ , we obtain from (2.15) that

$$\nu^{\lambda+1} \sin \theta \left| \mathcal{R}_{\nu}^{(\lambda)}(\theta) \right| \leq \frac{\lambda(\lambda-1)2^{3/2\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi}} \frac{\nu^{\lambda+1}}{(\nu+1)^{\lambda+1}} \left\{ |\cos \theta| + \frac{2}{3} \frac{\lambda+1}{\nu+1} \sin \theta + \frac{2^{2-\lambda} \Gamma(2\lambda-1)}{\Gamma(\lambda+1)} \frac{(\nu+1)^{\lambda+1}}{(\nu-\lambda+3)^{2\lambda-1}} |\cot \theta|^{\lambda-2} \left( |\cos \theta| + \frac{2}{3} \frac{2\lambda-1}{\nu-\lambda+3} \sin \theta \right) \right\}.$$

It is evident that

$$|\cos\theta| + \frac{2}{3}\frac{\lambda+1}{\nu+1}\sin\theta \le \frac{3\nu+2\lambda+5}{3(\nu+1)}, \quad |\cos\theta| + \frac{2}{3}\frac{2\lambda-1}{\nu-\lambda+3}\sin\theta \le \frac{3\nu+\lambda+7}{3(\nu-\lambda+3)}$$

We write

$$\frac{(\nu+1)^{\lambda+1}|\cot\theta|^{\lambda-2}}{(\nu-\lambda+3)^{2\lambda-1}} = \left(\frac{\nu+1}{\nu-\lambda+3}\right)^{\lambda+1} \left(\frac{\nu}{\nu-\lambda+3}\right)^{\lambda-2} \left(\frac{|\cot\theta|}{\nu}\right)^{\lambda-2}.$$

Using the inequality:  $\ln(1+z) \le z$  for z > -1 again, we derive

$$\left(\frac{\nu+1}{\nu-\lambda+3}\right)^{\lambda+1} = \exp\left((\lambda+1)\ln\left(1+\frac{\lambda-2}{\nu-\lambda+3}\right)\right) \le \exp\left(\frac{(\lambda-2)(\lambda+1)}{\nu-\lambda+3}\right),$$

and

$$\left(\frac{\nu}{\nu-\lambda+3}\right)^{\lambda-2} = \exp\left((\lambda-2)\ln\left(1+\frac{\lambda-3}{\nu-\lambda+3}\right)\right) \le \exp\left(\frac{(\lambda-3)(\lambda+1)}{\nu-\lambda+3}\right).$$

By (3.29), we have

$$\frac{1}{\nu\sin\theta} \le \frac{\pi}{2} \max\left\{\frac{1}{\nu\theta}, \frac{1}{\nu(\pi-\theta)}\right\} \le \frac{c\pi}{2},$$

which implies

$$\left(\frac{|\cot\theta|}{\nu}\right)^{\lambda-2} = |\cos\theta|^{\lambda-2} \left(\frac{1}{\nu\sin\theta}\right)^{\lambda-2} \le \left(\frac{c\pi}{2}\right)^{\lambda-2}.$$

We therefore derive from the above  $B_{\nu}^{(\lambda)}$  in the second case.

(iii) From (2.16), we obtain

$$\begin{split} \nu^{\lambda+1} \sin \theta \, |\mathcal{R}_{\nu}^{(\lambda)}(\theta)| \\ &\leq \frac{2^{\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi}} \frac{\nu^{\lambda+1}}{(\nu+\lambda-1)^{\lambda+1}} \left\{ \left. 2(\lambda+1/2) |\cos \theta| + \frac{(\lambda+1)^2}{\nu+\lambda} \sin \theta \right. \\ &+ (\lambda+2)(\lambda+1) \Big( 1 + \frac{2}{\nu+\lambda-1} - \frac{\lambda(\lambda+2)}{(\nu+\lambda-1)^2} \Big) \right. \\ &\times \Big( |\cos \theta| + \frac{2}{3} \frac{\lambda+3}{\nu+\lambda-1} \sin \theta \Big) \left. \right\} \\ &\leq \frac{2^{\lambda} \Gamma(\lambda+1/2)}{\sqrt{\pi}} \frac{\nu^{\lambda+1}}{(\nu+\lambda-1)^{\lambda+1}} \left\{ \left. 2(\lambda+1/2) + \frac{(\lambda+1)^2}{\nu+\lambda} \right. \\ &+ (\lambda+2)(\lambda+1) \Big( 1 + \frac{2}{\nu+\lambda-1} - \frac{\lambda(\lambda+2)}{(\nu+\lambda-1)^2} \Big) \Big( 1 + \frac{2}{3} \frac{\lambda+3}{\nu+\lambda-1} \Big) \right. \right\} \end{split}$$

This yields (2.20), and then the proof is completed.

# 3.4. Proof of Theorem 2.2

For  $\theta \in (0, \pi)$ , we can derive the bounds (2.24)–(2.25) from (2.13) by multiplying  $\sin \theta$  and  $(\sin \theta)^{\lambda-1}$ , respectively.

In order to obtain the upper bounds uniform for both  $\nu$  and  $\theta$ , it is necessary to study the behaviors of  ${}^{r}G_{\nu}^{(\lambda)}(x)$  at  $x = \pm 1$  (i.e.,  $\theta = 0, \pi$ ). It is evident that by (2.1),  ${}^{r}G_{\nu}^{(\lambda)}(1) = 1$  for all  $\lambda > -1/2$  and  $\nu \ge 0$ . We now examine the behavior of the right GGF-Fs at x = -1. It is clear that if  $\nu = n \in \mathbb{N}_0$ , we have  ${}^{r}G_n^{(\lambda)}(-1) = (-1)^n {}^{r}G_n^{(\lambda)}(1) = (-1)^n$ . We now consider the case with  $\nu \notin \mathbb{N}_0$ . Note that for  $-1/2 < \lambda < 1/2$  (cf. [13, Prop. 2.3]):

$${}^{r}G_{\nu}^{(\lambda)}(-1) = \frac{\cos((\nu+\lambda)\pi)}{\cos(\lambda\pi)},\tag{3.44}$$

so  ${}^{r}G_{\nu}^{(\lambda)}(x)$  is continuous on [-1, 1]. However, for  $\lambda \ge 1/2$  and  $\nu \notin \mathbb{N}_{0}$ ,  ${}^{r}G_{\nu}^{(\lambda)}(x)$  is singular at x = -1. Indeed, according to [13, Prop. 2.3], we have

$$\lim_{x \to -1^+} \frac{{}^{r} G_{\nu}^{(1/2)}(x)}{\ln(1+x)} = \frac{\sin(\nu\pi)}{\pi}, \quad \nu \notin \mathbb{N}_0;$$
(3.45)

and for  $\lambda > 1/2$  and  $\nu \notin \mathbb{N}_0$ , we have

$$\lim_{x \to -1^+} \left(\frac{1+x}{2}\right)^{\lambda - 1/2} {}^{r} G_{\nu}^{(\lambda)}(x) = -\frac{\sin(\nu\pi)}{\pi} \frac{\Gamma(\lambda - 1/2)\Gamma(\lambda + 1/2)\Gamma(\nu + 1)}{\Gamma(\nu + 2\lambda)} \coloneqq \mathcal{Q}_{\nu}^{(\lambda)}.$$
(3.46)

We remark that (3.46) also holds for  $\nu = n \in \mathbb{N}_0$ , as  $\mathcal{Q}_n^{(\lambda)} = 0$ .

We now consider the case with  $\theta = 0$ . As  ${}^{r}G_{\nu}^{(\lambda)}(1) = 1$ , taking the limit  $\theta \to 0$ , and find readily that the above bounds hold (note:  $\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(0) = 0$ , but  $\widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta) > 0$  in (2.24)–(2.25)).

It remains to consider  $\theta \to \pi^-$ , i.e.,  $x \to -1^+$ . Apparently, we have  $\sin \theta = \sqrt{1 - x^2}$ . As a direct consequence of (3.44)–(3.45), we have that for  $0 < \lambda \le 1/2$ ,

$$\lim_{\theta \to \pi^{-}} \widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = \lim_{\theta \to \pi^{-}} \left\{ (\sin \theta)^{\lambda + 1} {}^{r} G_{\nu}^{(\lambda)}(\cos \theta) \right\}$$
  
= 
$$\lim_{x \to -1^{+}} (1 - x^{2})^{(\lambda + 1)/2} {}^{r} G_{\nu}^{(\lambda)}(x) = 0.$$
 (3.47)

Similarly, by (3.46), we have that for  $1/2 < \lambda < 2$ ,

$$\lim_{\theta \to \pi^{-}} \widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = \lim_{x \to -1^{+}} \left\{ (1 - x^{2})^{1 - \lambda/2} (1 - x^{2})^{\lambda - 1/2} {}^{r} G_{\nu}^{(\lambda)}(x) \right\} = 0.$$
(3.48)

For  $\lambda \ge 2$ , we find from (3.46) that

$$\lim_{\theta \to \pi^{-}} \widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta) = \lim_{\theta \to \pi^{-}} \left\{ (\sin \theta)^{2\lambda - 1} \, {}^{r} G_{\nu}^{(\lambda)}(\cos \theta) \right\} = \lim_{x \to -1^{+}} (1 - x^{2})^{\lambda - 1/2} \, {}^{r} G_{\nu}^{(\lambda)}(x)$$

$$= 2^{2\lambda - 1} \, \mathcal{Q}_{\nu}^{(\lambda)}.$$
(3.49)

(i) For  $0 < \lambda \le 2$ ,  $\nu + \lambda > 1$  and  $\nu > 0$ , we find from (2.14) that

$$\lim_{\theta \to \pi^{-}} \widetilde{S}_{\nu}^{(\lambda)}(\theta) = \frac{\lambda |\lambda - 1| 2^{\lambda} \Gamma(\lambda + 1/2)}{\sqrt{\pi} (\nu + \lambda - 1)^{\lambda + 1}}.$$
(3.50)

Thus, in this case, it is evident that by (3.47)–(3.48) and (3.50), (2.24) holds for  $0 < \lambda < 2$ . For  $\lambda = 2$ , we obtain from (3.49)–(3.50) that

$$\lim_{\theta \to \pi^{-}} \widetilde{\mathcal{R}}_{\nu}^{(2)}(\theta) = 2^{3} \mathcal{Q}_{\nu}^{(2)} = -\frac{3\sin(\nu\pi)}{(\nu+1)(\nu+2)(\nu+3)}, \quad \lim_{\theta \to \pi^{-}} \widetilde{\mathcal{S}}_{\nu}^{(2)}(\theta) = \frac{6}{(\nu+1)^{3}}.$$

Hence, (2.24) holds for  $\lambda = 2$ .

(ii) For  $\lambda > 2$ ,  $\nu > \lambda - 3$  and  $\nu > 0$ , we obtain from (2.15) that

$$\lim_{\theta \to \pi^{-}} \widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta) = \frac{2^{2+\lambda/2} \Gamma(\lambda+1/2) \Gamma(2\lambda-1)}{\sqrt{\pi}(\nu-\lambda+3)^{2\lambda-1} \Gamma(\lambda-1)} = \frac{\lambda 2^{5\lambda/2} \Gamma(\lambda-1/2) \Gamma(\lambda+1/2)}{\pi(\nu-\lambda+3)^{2\lambda-1}}, \quad (3.51)$$

where we used the identity (cf. [1, (6.1.18)]):

 $\Gamma(2z) = \pi^{-1/2} 2^{2z-1} \Gamma(z) \Gamma(z+1/2).$ 

Using the inequality (cf. [17, (5.6.7)]): for  $b - a \ge 1$ ,  $a \ge 0$ , and z > 0,

$$\frac{\Gamma(z+a)}{\Gamma(z+b)} \le z^{a-b},$$

we get

$$\frac{\Gamma(\nu+1)}{\Gamma(\nu+2\lambda)} = \frac{\Gamma\left((\nu-\lambda+3)+(\lambda-2)\right)}{\Gamma\left((\nu-\lambda+3)+(3\lambda-3)\right)} \le (\nu-\lambda+3)^{1-2\lambda} \le \frac{\lambda 2^{\lambda/2+1}}{(\nu-\lambda+3)^{2\lambda-1}}.$$
 (3.52)

Thus, from (3.46) and (3.51)–(3.52), we derive that for  $\lambda \ge 2$ ,

$$\lim_{\theta \to \pi^{-}} |\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)| = 2^{2\lambda - 1} |\mathcal{Q}_{\nu}^{(\lambda)}| \leq \lim_{\theta \to \pi^{-}} \widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta).$$

(iii) For  $-\frac{1}{2} < \lambda < 0$  and  $\nu \geq 2$ , as  ${}^{r}G_{\nu}^{(\lambda)}(x)$  is continuous on [-1, 1], both  $\widetilde{\mathcal{R}}_{\nu}^{(\lambda)}(\theta)$  and  $\widetilde{\mathcal{S}}_{\nu}^{(\lambda)}(\theta)$  are continuous on  $[0, \pi]$ .

Therefore, the bounds in Theorem 2.2 are valid for all cases.

# 4. Some relevant properties of GGF-Fs

The GGF-Fs enjoy a rich collection of properties particularly in the fractional calculus framework. In this section, we present assorted properties of GGF-Fs, and most of them follow directly from the properties of the hypergeometric functions. These can provide a better picture of this family of very useful special functions.

Recall the definition of the right-sided Riemann–Liouville fractional derivative of order s > 0 (cf. [19]):

$${}_{x}^{R}D_{1}^{s}u(x) = (-1)^{k}D^{k}\left\{{}_{x}I_{1}^{k-s}u\right\}(x), \quad s \in [k-1,k),$$
(4.1)

where  $D^k$  with  $k \in \mathbb{N}$  is the ordinary kth derivative, and  ${}_xI_1^{\mu}$  is the RL fractional derivative operator defined in (2.8). We have the explicit formulas (cf. [19]): for real  $\eta > -1$  and s > 0,

$${}_{x}I_{1}^{s}(1-x)^{\eta} = \frac{\Gamma(\eta+1)}{\Gamma(\eta+s+1)}(1-x)^{\eta+s}; \quad {}_{x}^{R}D_{1}^{s}(1-x)^{\eta} = \frac{\Gamma(\eta+1)}{\Gamma(\eta-s+1)}(1-x)^{\eta-s}.$$
(4.2)

**Proposition 4.1** (See [13, Thm. 3.1]). For real  $\lambda > s - 1/2$ , real  $\nu \ge 0$  and  $x \in (-1, 1)$ ,

$${}^{R}_{x}D_{1}^{s}\left\{(1-x^{2})^{\lambda-1/2} \, {}^{r}G_{\nu}^{(\lambda)}(x)\right\} = \frac{2^{s} \, \Gamma(\lambda+1/2)}{\Gamma(\lambda-s+1/2)} \, (1-x^{2})^{\lambda-s-1/2} \, {}^{r}G_{\nu+s}^{(\lambda-s)}(x). \tag{4.3}$$

Note that we just list the properties for the right GGF-F  ${}^{r}G_{\nu}^{(\lambda)}(x)$ , but similar formulas are valid for the left GGF-F  ${}^{l}G_{\nu}^{(\lambda)}(x)$  (cf. (2.2)) under the left RL fractional derivative (cf. [13]).

As a generalization of Gegenbauer polynomials, the GGF-Fs satisfy the following fractional Rodrigues' formula.

**Proposition 4.2.** For real  $\lambda > -1/2$  and real  $\nu \ge 0$ , the GGF-Fs defined in (2.1) satisfy

$${}^{r}G_{\nu}^{(\lambda)}(x) = \frac{\Gamma(\lambda+1/2)}{2^{\nu}\,\Gamma(\nu+\lambda+1/2)}(1-x^{2})^{-\lambda+1/2} {}^{R}_{x}D_{1}^{\nu}\big\{(1-x^{2})^{\nu+\lambda-1/2}\big\}.$$
(4.4)

**Proof.** Substituting  $\nu$ ,  $\lambda$ , s in (4.3) by 0,  $\nu + \lambda$ ,  $\nu$ , respectively, yields

$${}_{x}^{R}D_{1}^{\nu}\left\{(1-x^{2})^{\nu+\lambda-1/2}\right\} = \frac{2^{\nu}\Gamma(\nu+\lambda+1/2)}{\Gamma(\lambda+1/2)}(1-x^{2})^{\lambda-1/2}G_{\nu}^{(\lambda)}(x),$$

which implies (4.4).

**Remark 4.1.** Mirevski et al. [15, Definition 9] defined the (generalized or) *g*-Jacobi function through the (fractional) Rodrigues' formula and derived an equivalent representation in terms of the hypergeometric function (cf. [15, Thm. 12]). However, we point out that the left RL fractional derivative operator  ${}_{0}^{R}D_{x}^{\nu}$  therein should be replaced by the right RL fractional derivative operator  ${}_{x}^{R}D_{1}^{\nu}$  as in (4.4). Then the flaws in the derivation of [15, Thm. 12] can be fixed accordingly.

**Proposition 4.3.** For real  $\lambda > -1/2$  and real  $\nu \ge 0$ , the GGF-Fs have the series representation:

$${}^{r}G_{\nu}^{(\lambda)}(x) = \frac{\Gamma(\lambda+1/2)\Gamma(\nu+1)}{2^{\nu}\Gamma(\nu+\lambda+1/2)} \sum_{k=0}^{\infty} \binom{\nu+\lambda-1/2}{\nu-k} \binom{\nu+\lambda-1/2}{k} (x-1)^{k}(1+x)^{\nu-k}.$$

**Proof.** Using the fractional Leibniz rule (cf. [18, (2.202)]), we obtain from (4.2) that

$${}^{R}_{x} D^{\nu}_{1} \{ (1-x^{2})^{\nu+\lambda-1/2} \} = \sum_{k=0}^{\infty} {\binom{\nu}{k}}^{R}_{x} D^{\nu-k}_{1} (1-x)^{\nu+\lambda-1/2} (-1)^{k} D^{k} (1+x)^{\nu+\lambda-1/2}$$
$$= (1-x^{2})^{\lambda-1/2} \sum_{k=0}^{\infty} {\binom{\nu}{k}} \frac{\Gamma^{2}(\nu+\lambda+1/2)(x-1)^{k}(1+x)^{\nu-k}}{\Gamma(k+\lambda+1/2)\Gamma(\nu-k+\lambda+1/2)}.$$

Recall the definition of the binomial coefficient

$$\binom{\nu}{k} = \frac{\Gamma(\nu+1)}{\Gamma(\nu-k+1)\Gamma(k+1)}.$$

Thus, we have

$$\binom{\nu+\lambda-1/2}{\nu-k} = \frac{\Gamma(\nu+\lambda+1/2)}{\Gamma(k+\lambda+1/2)\Gamma(\nu-k+1)},$$
$$\binom{\nu+\lambda-1/2}{k} = \frac{\Gamma(\nu+\lambda+1/2)}{\Gamma(\nu-k+\lambda+1/2)\Gamma(k+1)}.$$

Then the desired representation follows from the above.  $\Box$ 

**Remark 4.2.** Alternatively, we can derive the above identity from (2.1), (2.3) and the Pfaff's formula (cf. [2, Theorem 2.2.5]):

$$_{2}F_{1}(a, b; c; z) = (1 - z)^{-a} {}_{2}F_{1}(a, c - b; c; z/(1 - z)).$$

We next present some recurrence relations that generalize the corresponding formulas for the Gegenbauer polynomials.

**Proposition 4.4.** For real  $\lambda > -1/2$ , the GGF-Fs satisfy the recurrence formulas

$$(\nu + 2\lambda)^{r} G_{\nu+1}^{(\lambda)}(x) = 2(\nu + \lambda) x^{r} G_{\nu}^{(\lambda)}(x) - \nu^{r} G_{\nu-1}^{(\lambda)}(x), \quad \nu \ge 1,$$
(4.5)

and

$${}^{r}G_{\nu}^{(\lambda)}(x) = x \, {}^{r}G_{\nu-1}^{(\lambda+1)}(x) - \frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)}(1-x^{2}) \, {}^{r}G_{\nu-2}^{(\lambda+2)}(x), \quad \nu \ge 2.$$
(4.6)

**Proof.** Recall the formula (cf. [2, (2.5.15)]):

$$2b(c-a)(b-a-1){}_{2}F_{1}(a-1,b+1;c;z) -((1-2z)(b-a-1){}_{3}+(b-a)(b+a-1)(2c-b-a-1)){}_{2}F_{1}(a,b;c;z)$$
(4.7)  
$$-2a(b-c)(b-a+1){}_{2}F_{1}(a+1,b-1;c;z) = 0.$$

Substituting a, b, c and z in (4.7) by  $-\nu$ ,  $\nu + 2\lambda$ ,  $\lambda + 1/2$  and (1 - x)/2, respectively, and using (2.1), we obtain

$$2(\nu + 2\lambda)(\nu + \lambda + 1/2)(2\nu + 2\lambda - 1)^{r} G_{\nu+1}^{(\lambda)}(x) - (2\nu + 2\lambda - 1)_{3} x^{r} G_{\nu}^{(\lambda)}(x) + 2\nu(\nu + \lambda + 1/2)(2\nu + 2\lambda - 1)^{r} G_{\nu-1}^{(\lambda)}(x) = 0,$$
(4.8)

which implies (4.5).

Recall (cf. [2, (2.5.2)])

$$z(1-z)\frac{(a+1)(b+1)}{c(c+1)}{}_{2}F_{1}(a+2,b+2;c+2;z) + \frac{(c-(a+b+1)z)}{c}{}_{2}F_{1}(a+1,b+1;c+1;z) - {}_{2}F_{1}(a,b;c;z) = 0.$$
(4.9)

Substituting a, b, c and z in (4.9) by  $-\nu$ ,  $\nu + 2\lambda$ ,  $\lambda + 1/2$  and (1 - x)/2, respectively, leads to

$${}^{r}G_{\nu}^{(\lambda)}(x) = x \, {}^{r}G_{\nu-1}^{(\lambda+1)}(x) - \frac{(\nu-1)(\nu+2\lambda+1)}{4(\lambda+1/2)(\lambda+3/2)}(1-x^{2}) \, {}^{r}G_{\nu-2}^{(\lambda+2)}(x).$$

This completes the proof.  $\Box$ 

**Proposition 4.5.** For real  $\lambda > -1/2$  and real  $\nu \ge 0$ , the GGF-Fs satisfy the Sturm–Liouville equation

$$(1 - x^2)({}^{r}G_{\nu}^{(\lambda)}(x))'' - (2\lambda + 1)x\,({}^{r}G_{\nu}^{(\lambda)}(x))' + \nu(\nu + 2\lambda)\,{}^{r}G_{\nu}^{(\lambda)}(x) = 0, \tag{4.10}$$

or equivalently,

$$\left((1-x^2)^{\lambda+1/2} \left({}^r G_{\nu}^{(\lambda)}(x)\right)'\right)' + \nu(\nu+2\lambda)(1-x^2)^{\lambda-1/2} \, {}^r G_{\nu}^{(\lambda)}(x) = 0.$$
(4.11)

**Proof.** Note that  $F := {}_{2}F_{1}(a, b; c; z)$  satisfies the second-order equation (cf. [2, P. 94]):

$$z(1-z)F'' + (c - (a+b+1)z)F' - abF = 0.$$
(4.12)

Substituting *a*, *b*, *c* and *z* in (4.12) by  $-\nu$ ,  $\nu + 2\lambda$ ,  $\lambda + 1/2$  and (1 - x)/2, respectively, we derive (4.10) from (2.1).  $\Box$ 

Similar to the Gegenbauer polynomials, we have the following derivative relations.

**Proposition 4.6.** For real  $v \ge k \in \mathbb{N}$ , we have

$$\frac{d^k}{dx^k} {}^r G_{\nu}^{(\lambda)}(x) = (-1)^k \frac{(-\nu)_k (\nu + 2\lambda)_k}{2^k (\lambda + 1/2)_k} {}^r G_{\nu-k}^{(\lambda+k)}(x).$$
(4.13)

In particular, if k = 1, we have

$$\frac{d}{dx}^{r}G_{\nu}^{(\lambda)}(x) = \frac{\nu(\nu+2\lambda)}{2\lambda+1}^{r}G_{\nu-1}^{(\lambda+1)}(x), \quad \nu \ge 1.$$
(4.14)

**Proof.** The formula (4.13) is derived directly from the identity (cf. [17, (15.5.2)]):

$$\frac{d^k}{dz^k} {}_2F_1(a,b;c;z) = \frac{(a)_k(b)_k}{(c)_k} {}_2F_1(a+k,b+k;c+k;z),$$
(4.15)

and (2.1).  $\Box$ 

For completeness, we quote the following estimates, which were very useful in the error analysis in [13].

**Proposition 4.7** (See [13, Thms 2.1-2.2]). For  $0 < \lambda < 1$  and real  $\nu \ge 0$ , we have

$$\max_{|x| \le 1} \{ (1 - x^2)^{\lambda/2} \left|^r G_{\nu}^{(\lambda)}(x) \right| \} \le \varrho_{\nu}^{(\lambda)}, \tag{4.16}$$

where

$$\rho_{\nu}^{(\lambda)} = \frac{\Gamma(\lambda+1/2)}{\sqrt{\pi}} \left( \frac{\cos^2(\pi\nu/2)\Gamma^2(\nu/2+1/2)}{\Gamma^2((\nu+1)/2+\lambda)} + \frac{4\sin^2(\pi\nu/2)}{\nu^2+2\lambda\nu+\lambda} \frac{\Gamma^2(\nu/2+1)}{\Gamma^2(\nu/2+\lambda)} \right)^{1/2}.$$
(4.17)

For  $\lambda \geq 1$  and real  $\nu \geq 0$ , we have

$$\max_{|x| \le 1} \left\{ (1 - x^2)^{\lambda - 1/2} \left| {}^r G_{\nu}^{(\lambda)}(x) \right| \right\} \le \kappa_{\nu}^{(\lambda)},\tag{4.18}$$

where

$$\kappa_{\nu}^{(\lambda)} = \frac{\Gamma(\lambda + 1/2)}{\sqrt{\pi}} \left( \frac{\cos^2(\pi \nu/2)\Gamma^2((\nu+1)/2)}{\Gamma^2((\nu+1)/2 + \lambda)} + \frac{4\sin^2(\pi \nu/2)}{2\lambda - 1 + \nu(\nu+2\lambda)} \frac{\Gamma^2(\nu/2 + 1)}{\Gamma^2(\nu/2 + \lambda)} \right)^{1/2}.$$

It is seen from Lemma 3.3 and its proof in Appendix B that the bound (4.16) is useful for the derivation of the main result (2.16) in Theorem 2.1. Moreover, the special case with  $\nu = n$  improves the existing bound related to the usual Gegenbauer polynomials as remarked blow.

**Remark 4.3.** Recall the bound of Gegenbauer polynomial of degree *n* (cf. [17, (18.14.7)]): for  $0 < \lambda < 1$  and integer  $n \ge 0$ ,

$$\max_{|x| \le 1} \left\{ (1 - x^2)^{\lambda/2} \left| G_n^{(\lambda)}(x) \right| \right\} \le \frac{2^{1-\lambda} \Gamma(2\lambda) \Gamma(n+1)}{\Gamma(\lambda) \Gamma(n+2\lambda)} (n+\lambda)^{\lambda-1}.$$
(4.19)

In fact, we can show that

$$\varrho_n^{(\lambda)} \le \frac{\Gamma(\lambda+1/2)}{\sqrt{\pi}} \frac{\Gamma((n+1)/2)}{\Gamma((n+1)/2+\lambda)} \le \frac{2^{1-\lambda}\Gamma(2\lambda)\Gamma(n+1)}{\Gamma(\lambda)\Gamma(n+2\lambda)} (n+\lambda)^{\lambda-1}.$$
(4.20)

Therefore, the bound (4.16) is sharper than that of (4.19). The inequalities in (4.20) can be proved by following the same argument as for Lemma 3.3 in Appendix B. Here, we omit the derivation.

# Appendix A. Proof of Lemma 3.1

We first show that

$$\frac{t^2}{4}\cos^2\theta \left(\cosh(t/2)\right)^{\frac{4}{3}} + \sin^2\theta (\cosh t)^{\frac{2}{3}} < |g(\theta, t)|^2 < \frac{t^2}{4}\cos^2\theta \cosh^4(t/2) + \sin^2\theta \cosh^2 t,$$
(A.1)

and

$$|\partial_t g(\theta, t)| \le \left(\frac{t}{3}\sin\theta + \frac{1}{2}|\cos\theta|\right)\cosh t.$$
(A.2)

It is clear that

$$|g(\theta, t)|^{2} = \frac{\cos^{2}\theta(\cosh t - 1)^{2} + \sin^{2}\theta\sinh^{2}t}{t^{2}}.$$
(A.3)

Recall the properties of hyperbolic functions (cf. [17, (4.32.1), (4.32.2), (4.35.20)]): for t > 0,

$$(\cosh t)^{\frac{1}{3}} < \frac{\sinh t}{t}; \quad \tanh t < t; \quad \sinh \frac{t}{2} = \left(\frac{\cosh t - 1}{2}\right)^{\frac{1}{2}}.$$
 (A.4)

Then we derive

$$(\cosh t)^{\frac{1}{3}} < \frac{\sinh t}{t} < \cosh t, \quad \forall t > 0, \tag{A.5}$$

and

$$\frac{1}{2} \left( \cosh(t/2) \right)^{\frac{2}{3}} < \frac{\cosh t - 1}{t^2} = \frac{1}{2} \left( \frac{\sinh(t/2)}{t/2} \right)^2 < \frac{1}{2} \cosh^2(t/2).$$
(A.6)

Thus we obtain (A.1) from (A.3) and (A.5)–(A.6) immediately.

Direct calculation from (3.1) yields

$$\partial_t g(\theta, t) = \frac{\cos \theta \left( t \sinh t - \cosh t + 1 \right) + i \sin \theta \left( t \cosh t - \sinh t \right)}{t^2},\tag{A.7}$$

and

$$\left|\partial_{t}g(\theta,t)\right|^{2} = \frac{\cos^{2}\theta \left(t\sinh t - \cosh t + 1\right)^{2} + \sin^{2}\theta \left(t\cosh t - \sinh t\right)^{2}}{t^{4}}.$$
 (A.8)

We next show that for t > 0,

$$\frac{t\cosh t - \sinh t}{t^3} < \frac{1}{3}\cosh t,\tag{A.9}$$

and

$$\frac{t\sinh t - \cosh t + 1}{t^2} < \frac{1}{2}\cosh t.$$
(A.10)

To prove (A.9), we denote  $h(t) := t^3 \cosh t - 3t \cosh t + 3 \sinh t$ . Then for t > 0,

$$h'(t) = t^{3} \sinh t + 3t(t \cosh t - \sinh t) > t^{3} \sinh t > 0,$$
(A.11)

where we used the property:  $t \cosh t > \sinh t$  (cf. (A.4)). Therefore, h(t) is strictly ascending, so for all t > 0,

$$h(t) = t^{3} \cosh t - 3t \cosh t + 3 \sinh t > h(0) = 0,$$

which implies (A.9). As

$$(t \sinh t - \cosh t + 1)' = t \cosh t > 0, \quad t > 0,$$

we have  $t \sinh t - \cosh t + 1 > 0$  for all t > 0. Denoting  $\hat{h}(t) := t^2 \cosh t - 2t \sinh t + 2 \cosh t - 2$ , we find for t > 0,

$$\hat{h}'(t) = t^2 \sinh t > 0$$
, so  $\hat{h}(t) > \hat{h}(0) = 0$ , (A.12)

which yields (A.10).

From (A.8)–(A.10), we obtain

$$|\partial_t g(\theta, t)|^2 \le \frac{1}{9} t^2 \sin^2 \theta \cosh^2 t + \frac{1}{4} \cos^2 \theta \cosh^2 t, \qquad (A.13)$$

which leads to (A.2).

Now, we are ready to derive (3.2)–(3.3). Using the mean-value theorem for the real part and imaginary part of  $f^{(\lambda)}(\theta, t)$ , respectively, we obtain

$$f^{(\lambda)}(\theta,t) = \frac{g^{\lambda-1}(\theta,t) - g^{\lambda-1}(\theta,0)}{t} = \Re \{\partial_t g^{\lambda-1}(\theta,\xi_1)\} + i\Im \{\partial_t g^{\lambda-1}(\theta,\xi_2)\}, \quad (A.14)$$

for  $\xi_i = \xi_i(t) \in (0, t)$ , i = 1, 2, and  $\theta \in (0, \pi)$ . Hence, we have

$$|f^{(\lambda)}(\theta,t)| \le 2 \sup_{0<\xi< t} |\partial_t g^{\lambda-1}(\theta,\xi)| = 2|\lambda-1| \sup_{0<\xi< t} \{ |g(\theta,\xi)|^{\lambda-2} |\partial_t g(\theta,\xi)| \}.$$
(A.15)

We now estimate its upper bound. From (A.2), we obtain that for  $\xi \in (0, t)$  and  $\theta \in (0, \pi)$ ,

$$|\partial_t g(\theta,\xi)| \le \left(\frac{\xi}{3}\sin\theta + \frac{1}{2}|\cos\theta|\right)\cosh\xi \le \left(\frac{t}{3}\sin\theta + \frac{1}{2}|\cos\theta|\right)e^t.$$
(A.16)

It remains to estimate the upper bound of  $|g(\theta, \xi)|^{\lambda-2}$ . We proceed with two cases. (i) For  $0 < \lambda \le 2$ , we obtain from the lower bound g in (A.1) that for  $0 < \xi < t$ ,

$$|g(\theta,\xi)|^{\lambda-2} \le \left(\frac{1}{4}\cos^2\theta \cosh^{4/3}(\xi/2)\xi^2 + \sin^2\theta \cosh^{2/3}\xi\right)^{\lambda/2-1} \le \sin^{\lambda-2}\theta, \qquad (A.17)$$

where we used the fact the function in  $\xi$  is strictly decreasing, since  $\lambda/2 - 1 < 0$ . Thus, we obtain (3.2) from (A.15)–(A.17).

(ii) For  $\lambda > 2$ , we obtain from the upper bound of g in (A.1) that

$$\begin{split} |g(\theta,\xi)|^{\lambda-2} &\leq \left(\frac{1}{4}\cos^2\theta\cosh^4(\xi/2)\xi^2 + \sin^2\theta\cosh^2\xi\right)^{\lambda/2-1} \\ &\leq \left(\frac{1}{4}\cos^2\theta\cosh^4(t/2)t^2 + \sin^2\theta\cosh^2t\right)^{\lambda/2-1} \\ &\leq \left(\max\left\{\frac{1}{2}\cos^2\theta\cosh^4(t/2)t^2, 2\sin^2\theta\cosh^2t\right\}\right)^{\lambda/2-1} \\ &\leq \left(\max\left\{\frac{1}{2}t^2\cos^2\theta\cosh^4t, 2\sin^2\theta\cosh^2t\right\}\right)^{\lambda/2-1} \\ &\leq 2^{1-\lambda/2}(\cos\theta)^{\lambda-2}(\cosh t)^{2\lambda-4}t^{\lambda-2} + 2^{\lambda/2-1}(\sin\theta)^{\lambda-2}(\cosh t)^{\lambda-2} \\ &\leq 2^{1-\lambda/2}(\cos\theta)^{\lambda-2}e^{2(\lambda-2)t}t^{\lambda-2} + 2^{\lambda/2-1}(\sin\theta)^{\lambda-2}e^{(\lambda-2)t} \\ &= 2^{\lambda/2-1}(\sin\theta)^{\lambda-2}e^{(\lambda-2)t}\left(1 + \frac{|\cot\theta|^{\lambda-2}}{2^{\lambda-2}}t^{\lambda-2}e^{(\lambda-2)t}\right). \end{split}$$
(A.18)

Therefore, we obtain (3.3) from (A.15)–(A.16) and (A.18).

# Appendix B. Proof of Lemma 3.3

We first show that for  $0 < \lambda < 1$  and real  $\nu \ge 0$ , we have

$$\max_{|x| \le 1} \left\{ (1 - x^2)^{\lambda/2} \left|^r G_{\nu}^{(\lambda)}(x) \right| \right\} \le \frac{\Gamma(\lambda + 1/2)}{\sqrt{\pi}} \frac{\Gamma((\nu + 1)/2)}{\Gamma((\nu + 1)/2 + \lambda)}.$$
(B.1)

In view of (4.16)–(4.17), it suffices to prove

$$\varrho_{\nu}^{(\lambda)} \leq \frac{\Gamma(\lambda+1/2)}{\sqrt{\pi}} \frac{\Gamma((\nu+1)/2)}{\Gamma((\nu+1)/2+\lambda)}.$$
(B.2)

Indeed, the inequality (B.2) holds, if we can show

$$\frac{4}{\nu^2 + 2\lambda\nu + \lambda} \frac{\Gamma^2(\nu/2 + 1)}{\Gamma^2(\nu/2 + \lambda)} \le \frac{\Gamma^2((\nu + 1)/2)}{\Gamma^2((\nu + 1)/2 + \lambda)},$$
(B.3)

thanks to the expression of  $\rho_{\nu}^{(\lambda)}$  in (4.17).

Next we prove (B.3). For notational convenience, we denote

$$\Upsilon_{\lambda}(\nu) = \frac{4}{\nu^2 + 2\lambda\nu + \lambda} \frac{\Gamma^2(\nu/2 + 1)}{\Gamma^2(\nu/2 + \lambda)} \frac{\Gamma^2((\nu + 1)/2 + \lambda)}{\Gamma^2((\nu + 1)/2)},$$
(B.4)

and

$$\Phi_{\lambda}(\nu) = \frac{\Upsilon_{\lambda}(\nu)}{\Upsilon_{\lambda}(\nu+2)} = \frac{(\nu+2)^2 + 2\lambda(\nu+2) + \lambda}{\nu^2 + 2\lambda\nu + \lambda} \frac{(\nu/2+\lambda)^2}{(\nu/2+1)^2} \frac{(\nu+1)^2/4}{((\nu+1)/2+\lambda)^2}, \quad (B.5)$$

where we used the property:  $\Gamma(z + 1) = z\Gamma(z)$ . Then direct calculation leads to

$$\begin{split} \frac{\Phi'_{\lambda}(\nu)}{\Phi_{\lambda}(\nu)} &= (\ln \Phi_{\lambda}(\nu))' = \frac{2(\nu+2)+2\lambda}{(\nu+2)^2+2\lambda(\nu+2)+\lambda} - \frac{2\nu+2\lambda}{\nu^2+2\lambda\nu+\lambda} \\ &+ \frac{2}{\nu+2\lambda} - \frac{2}{\nu+2} + \frac{2}{\nu+1} - \frac{2}{\nu+1+2\lambda} \\ &= -4 \frac{(\nu+\lambda+1)^2 - 1 - \lambda + \lambda^2}{((\nu+\lambda)^2+\lambda-\lambda^2)((\nu+\lambda+2)^2+\lambda-\lambda^2)} + \frac{2}{(\nu+2\lambda)(\nu+2\lambda+1)} \\ &+ \frac{2}{(\nu+1)(\nu+2)} \\ &= \frac{4\lambda(1-\lambda)(5\nu^2+10(1+\lambda)\nu+6\lambda^2+9\lambda+4)}{(\nu+1)(\nu+2)(\nu+2\lambda)(\nu+2\lambda+1)((\nu+\lambda)^2+\lambda-\lambda^2)((\nu+\lambda+2)^2+\lambda-\lambda^2)} \\ &\geq 0. \end{split}$$

Thus,  $\Phi_{\lambda}(\nu)$  is an increasing function in  $\nu$ , which, together with (B.5), yields

$$\frac{\Upsilon_{\lambda}(\nu)}{\Upsilon_{\lambda}(\nu+2)} = \varPhi_{\lambda}(\nu) \le \lim_{\nu \to \infty} \varPhi_{\lambda}(\nu) = 1.$$
(B.6)

This implies

$$\Upsilon_{\lambda}(\nu) \leq \Upsilon_{\lambda}(\nu+2) \leq \Upsilon_{\lambda}(\nu+4) \leq \cdots \leq \Upsilon_{\lambda}(\nu+2n),$$
(B.7)

for any positive integer n. Consequently, we have

$$\frac{4}{\nu^2 + 2\lambda\nu + \lambda} \frac{\Gamma^2(\nu/2+1)}{\Gamma^2(\nu/2+\lambda)} \frac{\Gamma^2((\nu+1)/2+\lambda)}{\Gamma^2((\nu+1)/2)} = \Upsilon_{\lambda}(\nu) \le \lim_{n \to \infty} \Upsilon_{\lambda}(\nu+2n) = 1, \quad (B.8)$$

where we used the property (2.23) for finding the limit value. From (B.8), we obtain (B.3), which implies (B.1)–(B.2).

The second step is to show that for  $0 < \lambda < 1$  and  $\nu \ge 0$ ,

$$\frac{\Gamma((\nu+1)/2)}{\Gamma((\nu+1)/2+\lambda)} \le \left((\nu+\lambda)/2\right)^{-\lambda},\tag{B.9}$$

as it ensures

$$\frac{\Gamma(\lambda+1/2)}{\sqrt{\pi}} \frac{\Gamma((\nu+1)/2)}{\Gamma((\nu+1)/2+\lambda)} \leq \frac{2^{\lambda}\Gamma(\lambda+1/2)}{\sqrt{\pi}(\nu+\lambda)^{\lambda}}$$

As such, if (B.9) holds, then (3.32) is a direct consequence of (B.1). To derive (B.9), we denote

$$\widehat{\Upsilon}_{\lambda}(\nu) = \frac{(\nu/2 + \lambda/2)^{\lambda} \Gamma((\nu+1)/2)}{\Gamma((\nu+1)/2 + \lambda)},$$

and

$$\widehat{\varPhi}_{\lambda}(\nu) = \frac{\widehat{\Upsilon}_{\lambda}(\nu)}{\widehat{\Upsilon}_{\lambda}(\nu+2)} = \frac{(\nu/2 + \lambda/2)^{\lambda}}{(\nu/2 + 1 + \lambda/2)^{\lambda}} \frac{(\nu+1)/2 + \lambda}{(\nu+1)/2}.$$

One verifies readily that

$$\begin{split} \widehat{\varPhi}'_{\lambda}(\nu) &= (\ln \widehat{\varPhi}_{\lambda}(\nu))' = \frac{\lambda}{\nu+\lambda} - \frac{\lambda}{\nu+2+\lambda} + \frac{1}{\nu+1+2\lambda} - \frac{1}{\nu+1} \\ &= \frac{2\lambda(1-\lambda^2)}{(\nu+\lambda)(\nu+1)(\nu+1+2\lambda)(\nu+2+\lambda)} \ge 0. \end{split}$$

so  $\widehat{\Phi}_{\lambda}(\nu)$  is an increasing function in  $\nu$ . Following the same lines as in the derivations in (B.6)–(B.8), we can derive (B.9). This competes the proof.

#### References

- M. Abramowitz, I.A. Stegun, Handbook of Mathematical Functions with Formulas, in: Graphs and Mathematical Tables, Dover, New York, 1972.
- [2] G.E. Andrews, R. Askey, R. Roy, Special Functions, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, 1999.
- [3] I. Babuška, B.Q. Guo, Optimal estimates for lower and upper bounds of approximation errors in the *p*-version of the finite element method in two dimensions, Numer. Math. 85 (2000) 219–255.
- [4] I. Babuška, B.Q. Guo, Direct and inverse approximation theorems for the *p*-version of the finite element method in the framework of weighted Besov spaces I: Approximability of functions in the weighted Besov spaces, SIAM J. Numer. Anal. 39 (5) (2001) 1512–1538.
- [5] I. Babuška, B.Q. Guo, Direct and inverse approximation theorems for the *p*-version of the finite element method in the framework of weighted Besov spacesm, Part II: Optimal rate of convergence of the *p*-version finite element solutions, Math. Models Methods Appl. Sci. 12 (5) (2002) 689–719.
- [6] I. Babuška, M. Suri, The hp-version of the finite element method with quasi-uniform meshes, RAIRO Model. Math. Anal. Numer. 21 (1987) 199–238.
- [7] S. Chen, J. Shen, L.L. Wang, Generalized Jacobi functions and their applications to fractional differential equations, Math. Comput. 85 (300) (2016) 1603–1638.
- [8] S. Farid Khwaja, A.B. Olde Daalhuis, Uniform asymptotic expansions for hypergeometric functions with large parameters, IV, Anal. Appl. 12 (6) (2014) 667–710.
- [9] W. Gui, I. Babuška, The h, p and h-p versions of the finite element method in 1 dimension, Part I: The error analysis of the *p*-version, Numer. Math. 49 (1986) 205–612.
- [10] B.Y. Guo, J. Shen, L.L. Wang, Generalized Jacobi polynomials/functions and their applications, Appl. Numer. Math. 59 (5) (2009) 1011–1028.
- [11] B.Y. Guo, L.L. Wang, Jacobi approximations in non-uniformly Jacobi-weighted Sobolev spaces, J. Approx. Theory 128 (1) (2004) 1–41.
- [12] D. Jones, Asymptotics of the hypergeometric function, Math. Methods Appl. Sci. 24 (2001) 369–389.
- [13] W.J. Liu, L.L. Wang, H.Y. Li, Optimal error estimates for Chebyshev approximations of functions with limited regularity in fractional Sobolev-type spaces, Math. Comput. 88 (320) (2019) 2857–2895.
- [14] D.S. Lubinsky, Asymptotics of orthogonal polynomials: Some old, some new, some identities, Acta Appl. Math. 61 (1) (2000) 207–256.
- [15] S.P. Mirevski, L. Boyadjiev, R. Scherer, On the Riemann-Liouville fractional calculus, *g*-Jacobi functions and *F*-Gauss functions, Appl. Math. Comput. 187 (2007) 315–325.
- [16] F.W.J. Olver, Asymptotics and Special Functions, Academic Press, New York, 1974.
- [17] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clark, NIST Handbook of Mathematical Functions, Cambridge University Press, New York, 2010.
- [18] I. Podlubny, Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications, in: Mathematics in Science and Engineering, vol. 198, Academic Press, Inc. San Diego, CA, 1999.
- [19] S.G. Samko, A.A. Kilbas, O.I. Marichev, Fractional Integrals and Derivatives, Theory and Applications, Gordan and Breach Science Publisher, New York, 1993.
- [20] C. Schwab, P- and hp-FEM: Theory and Application to Solid and Fluid Mechanics, Oxford University Press, New York, 1998.
- [21] J. Shen, T. Tang, L.L. Wang, Spectral Methods: Algorithms, Analysis and Applications, Springer-Verlag, New York, 2011.
- [22] G. Szegö, Orthogonal Polynomials, fourth ed., Amer. Math. Soc., Providence, RI, 1975.
- [23] N.M. Temme, Special Functions: An Introduction to the Classical Functions of Mathematical Physics, Wiley, New York, 1996.
- [24] L.N. Trefethen, Approximation Theory and Approximation Practice, SIAM, Philadelphia, 2013.
- [25] G. Watson, Asymptotic expansions of hypergeometric functions, Trans. Cambridge Philos. Soc. 22 (1918) 277–308.
- [26] R. Wong, Orthogonal polynomials and their asymptotic behavior, in: C. Dunkl, M. Ismail, R. Wong (Eds.), Proceeding of the International Workshop on Special Functions, World Scientific, 2000, pp. 409–422.
- [27] R. Wong, Asymptotics of orthogonal polynomials, Int. J. Numer. Anal. Model. 15 (1-2) (2018) 193-212.
- [28] M. Zayernouri, G.E. Karniadakis, Fractional Sturm-Liouville eigen-problems: Theory and numerical approximation, J. Comput. Phys. 252 (2013) 495–517.